

Effects of Exogenous Salicylic Acid on Morphological and Biochemical Characteristics of *Jatropha Curcas* L. Irrigated with Saline Water

*Abo El-Soud, I.H.¹ and Hokam, E.M.²

¹ Horticulture Departments, Fac. of Agric., Suez-Canal Univ., Ismailia, Egypt. 41522

² Soil and Water Departments, Fac. of Agric., Suez-Canal Univ., Ismailia, Egypt. 41522

* E-mail: Islamhassan2010@hotmail.com

Received on: 15/1/2017

Accepted for publication on: 19/1/2017

Abstract

Salinity is one of the major rate-limiting to crop productivity. A field study was conducted at two developmental stages of *Jatropha curcas* L. namely: the juvenile stage (1-2 years) and the early mature stage (3-4 years). Three levels of water salinity were prepared by diluting sea water (Suez Canal water) to give 2.3, 4.7 and 7.0 dSm⁻¹, in addition to fresh water (0.32dSm⁻¹). *Jatropha* plants were foliar sprayed with three levels of salicylic acid (0.0, 5.0 and 10 mM). On the juvenile stage, the obtained results indicated that the most effective treatment for vegetative growth characters such as total leaf area (TLA); leaf relative water content (RWC_L); relative growth rate (RGR) and leaf dry weight (LDW) was performed when *Jatropha* plants were irrigated with fresh water, without significant difference, than that irrigated plants with either 2.3 or 4.7 dSm⁻¹. Application of 5.0 mM salicylic acid (SA) increased RGR, TLA and LDW for plants irrigated with saline water at 4.7 dSm⁻¹ as compared to either 0.0 mM or 10 mM SA.

On the early mature stage, application of 5.0 mM SA had a positive effect on both vegetative characters and seed yield for 4.7dSm⁻¹ salinity treatments as compared to control treatment. On the other hand, values of RWC_L indicated that there was no water deficit in leaves of treatments irrigated with saline water or control. This indicates that application of SA may minimize salinity stress. Foliar application of SA effectively increased chlorophyll. Generally, spraying *jatropha* shrubs with SA had a positive effect in reducing the negative impact of water salinity.

Keywords: *Jatropha curcas*; *physic nut*; *physic nut oil*; *Salicylic Acid*; *sea water*, *irrigation*; *biofuel plant*.

Introduction

The desert in Egypt occupies 94% of total area. A small number of crops can be cultivated in the deserts. Salt stresses either in soils or irrigation water are the major abiotic stresses that limit crop production in deserts (Allakhverdiev *et al.*, 2000). Salt stress alters many different physiological and biochemical responses in plants, thereby affecting almost all plant processes (Iqbal *et al.*, 2006). The important effects of

salinity are reductions in plant growth parameters, such as decreased leaf area, leaf length and root and shoot dry weight (Saleh, 2012). Besides, irrigation with high-quality water has become along beggarly in many areas (Niu *et al.*, 2012).

Salicylic acid (SA) has been shown as an important signal molecule for modulating plant responses to abiotic stresses (Khan *et al.*, 2015). The effects of SA on plant physiological processes varies depending on

species, developmental stage, SA concentration and the environmental conditions (Shraiy and Hegazi, 2009). The positive effects of SA have been documented in inducing salinity tolerance of several crops (Khan *et al.*, 2015). Data obtained by Najafian *et al.*, (2009) showed that plants treated with SA under salt stress had greater shoot and root dry weights, as well as SA increased photosynthetic rate and nutrient uptake.

The demand for biofuel has been rapidly growing in the last decade, global production of biofuel more than doubled (Licht, 2009). The potential resources for second-generation for biofuels production are *Jatropha* and algae. On the other hand, water scarcity is one of the major constraints of future potential production of second generation of biofuels (Hoekstra and Chapagain 2007). It is important for the development countries "such as Egypt" to start production of biodiesel to achieve the objectives of emission standards, and benefit from degraded lands and encourage the growth of the renewable energy industry (Croker, 2013).

Jatropha curcas (family Euphorbiaceae), a deciduous shrub, can be grown in semi-arid conditions, and marginal soils without large invest-

ments in inputs (Jongschaap *et al.*, 2007). *Jatropha* is commonly used for several purposes such as a natural hedge or fence-row, oil for soap production and cosmetics. Various parts of the plant have medicinal value; and can be used as an organic "green manure" fertilizer (Jewitt *et al.*, 2009). About Five million hectares of *Jatropha* crop are being established on a worldwide scale, scattered across a vast number of countries. On the other hand, 700 feddan is now cultivated in Egypt, in Ismailia, Suez, Luxor and Giza governorates. The objective of this study was to investigate the effects of foliar spraying *J. Curcas* L. plants with salicylic acid on growth and oil yield under irrigation with saline water.

Materials and Methods

A field study was conducted at the Horticulture farm, Suez-Canal Univ., Ismailia, Egypt, from 2012 to 2015 on sandy soil (Table 1). This study has been carried out at two stages of growth of *Jatropha* plants. The first is the juvenile stage from transplanting to the end of the second year (without bearing fruits). Meanwhile, the second is the early stage of maturity up to the fourth year where plants begin to produce fruits.

Table 1. Some physical and chemical properties of soil

Physical properties								
Coarse sand	Fine sand	Silt	Clay	Texture class	Field capacity			
81.0 %	15.3%	0.9%	2.8%	Sandy	0.13 cm ³ cm ⁻³			
Chemical properties								
Electrical conductivity, dS m ⁻¹	Organic matter %	N _{total} %	P %	K %	Ca ppm	Na ppm	Cl ppm	
0.98	0.39	0.25	.008	0.03	88	107.3	211.7	

Jatropha seeds were planted directly into the field in Mid of October, 2011 at 2.5 m x 2.5 m. Three levels of irrigation water salinity based on water electric conductivity (EC), were prepared by diluting sea water (from Suez Canal) to gave 2.3, 4.7 and 7.0 dSm⁻¹, in addition to the fresh water as control (0.32 dSm⁻¹). Chemical analysis of diluted sea water used in irrigation is presented in Table (2). In Mid March (2012), *Jatropha* seedlings were foliar sprayed

with three levels of SA as 0.0, 5.0 and 10 mM. SA was dissolved in small volume of ethanol and the solution was completed with water then Tween 20 was added to decrease the surface tension of the solution.

Plants sprayed with SA at four weeks interval, making six applications during the growth season. Plants were regular fertilized with complete fertilizers at the ratio of 6:10:6 and micronutrients. Fertilizers were applied every 2 weeks for 20 times.

Table 2. Chemical properties of the irrigation waters

Parameters	EC, dS m ⁻¹	pH	Soluble cations, meq l ⁻¹				Soluble anions, meq l ⁻¹			
			Ca ²⁺	Mg ²⁺	Na ⁺	K ⁺	Cl ⁻	HCO ₃ ⁻	SO ₄ ²⁻	SAR ^{**}
Fresh water	0.32	7.95	0.97	0.60	1.64	0.39	1.50	1.30	0.80	1.85
Diluted sea water*	7.00	7.67	6.00	9.00	57.5	1.50	49.0	3.40	21.6	21.0

*Sea water: Nile water(1 :10) **SAR: Sodium adsorption ratio

Relative growth rate (RGR), was calculated according to Antúnez (2001); total leaf area (TLA) according to Pérez-Harguindeguy *et al.*, (2013); leaf dry weight (LDW), and seed yield plant⁻¹. Concerning seed oil percentage, mature fruits were collected and air dried at room temperature. Then the seeds were crushed, and oil was extracted by hexane according to the method described in Bilal *et al.*, (2013). Relative water content (leaves) RWCL was calculated according to the method described by Villar-Salvador *et al.*, (2012). Total chlorophyll was determined and calculated according to the method described by Shibghatallah *et al.*, (2013).

At the end of the experiment during August, samples were taken from third leaf from the plant apex, cleaned with distilled water, dried at 70°C to constant weight and finally grind to determine N, P and K content. Total nitrogen percentage, was

determined by micro-Kjeldahl according to the method described by Jones (2001). Phosphorus percentage was estimated calorimetrically as described by Mazumder and Majumdar (2003). Potassium and sodium percentages were determined using the flame photometer according to Jones (2001).

Statistical Analyses:

In a factorial experiment, treatments were performed in a randomized complete blocks design with three replications and three plants in each replicate. Data were subjected to ANOVA using General Linear Model. Means of the interaction effects were compared using Duncan at 5% level of probability. Data were analyzed by SPSS program.

Results and Discussions

After 4 years of treatments (about 1450 days), the salinity induced modulations of growth were assessed by analyzing RGR for plant height, total leaf area and dry biomass of leaves (Tables 3 and 5). Concerning the main effect of salinity, there was no significant changes in RGR for plant height between control and 2.3 dSm⁻¹ and 4.7 dSm⁻¹ treated plants, respectively. The total leaf area was decreased significantly by 70.05% at 4.7 dSm⁻¹ compared to control (Table 5). The same conclusions were reported by Maggio *et al.*, (2000) who found that salinity effects on ion accumulation have been reported in *Salvadora persica* plants treated with salinity up to 200 mM NaCl.

Table (3) shows that most of the vegetative characters (Total leaf area (TLA); Relative water content of leaves (RWC_L); Height-Relative growth rate (height -RGR); leaf dry weight (LDW) were significantly affected by interaction between SA concentrations and water salinity in both years. Results indicated that the most effective treatment was found when *Jatropha* was irrigated with the first level of salinity as 2.3 dSm⁻¹ and sprayed with 5.0 mM SA, but without significant differences with treatments irrigated with either 0.32 or 4.70dSm⁻¹.

Generally, spraying plants with 5.0 mM SA positively affected the

growth characters at salinity level of 4.7 dSm⁻¹ compared with either 10.0 mM or without SA. Similar results were obtained by Dagar *et al.*, (2006) and Kotoky *et al.*, (2015). At the same effort, Fujimaki and Kikuchi (2010) reported that *Jatropha* is not more tolerance to salinity compared to other major crops such as soybean or wheat. According to Niu *et al.*, (2012) salinity stress and deficit irrigation significantly reduced the growth and leaf development of greenhouse grown *Jatropha* plants. They added that the degree of the foliar salt damage was increased with increasing the salinity of irrigation water.

It is evident from Table (3) that the RWC of *Jatropha* was affected by salicylic acid and salinity and their interactions. Concerning the main effect of salinity, regardless salicylic acid, the largest increase in RWC was recorded in the treatment irrigated with 7.0 dSm⁻¹ salinity. Meanwhile, in the case of spraying salicylic acid, regardless to the effect of salinity, spraying plants with 5.0 mM salicylic acid or without spraying salicylic acid, showed higher values of RWC without significant difference between them. RWC in leaves was 77.9% and 77.2% in control treatment while it was 86.3% and 88.4% in plants irrigated with 7.0 dSm⁻¹ in the first and second years, respectively (Table 3).

Table 3. Main and interactions effects of water salinity levels and SA concentrations on vegetative growth characters of *Jatropha* at juvenile stage (1st and 2nd years).

EC dSm ⁻¹	SA mM	TLA cm ²		RWC _L %		Height RGR cm month ⁻¹		LDW g plant ⁻¹	
		1 st year	2 nd year	1 st year	2 nd year	1 st year	2 nd year	1 st year	2 nd year
Main effects of water salinity									
0.32		135.7 ^a	329.6 ^b	77.9 ^b	77.2 ^b	1.4 ^a	0.41 ^b	0.38 ^d	0.97 ^c
2.3		126.4 ^a	378.8 ^a	76.8 ^b	77.2 ^b	1.4 ^a	0.41 ^b	0.57 ^b	1.8 ^a
4.7		126.4 ^a	340.1 ^b	81.2 ^{ab}	80.5 ^b	1.4 ^a	0.49 ^a	0.68 ^a	1.7 ^a
7.0		115.7 ^b	284.1 ^{bc}	86.3 ^a	88.4 ^a	1.4 ^a	0.26 ^c	0.51 ^c	1.40 ^b
Main effects of SA concentrations									
	0	112.4 ^c	377.5 ^b	82.8 ^a	84.0 ^a	1.3 ^b	0.36 ^c	0.40 ^b	1.55 ^a
	5	143.2 ^a	409.5 ^a	81.5 ^a	81.9 ^{ab}	1.4 ^{ab}	0.42 ^a	0.59 ^a	1.56 ^a
	10	122.6 ^b	212.5 ^c	77.4 ^a	76.7 ^b	1.5 ^a	0.39 ^b	0.55 ^a	1.32 ^b
Interaction effects of water salinity and SA concentrations									
0.32	0	104.0 ^c	437.2 ^a	85.1 ^{ab}	85.5 ^{ab}	1.14 ^d	0.47 ^c	0.43 ^e	1.79 ^c
	5	185.6 ^a	360.8 ^b	77.02 ^{abc}	76.4 ^{bc}	1.58 ^a	0.38 ^{de}	0.31 ^f	0.61 ^f
	10	117.6 ^{cde}	190.8 ^c	71.7 ^c	69.8 ^c	1.42 ^{ab}	0.38 ^{de}	0.39 ^e	0.52 ^f
2.3	0	128.0 ^{bcd}	448.7 ^a	83.26 ^{abc}	85.5 ^{ab}	1.5 ^{8a}	0.43 ^{cd}	0.41 ^e	1.48 ^d
	5	136.0 ^{bc}	477.7 ^a	75.34 ^{bc}	76.4 ^{bc}	1.06 ^d	0.45 ^c	0.72 ^b	2.10 ^{ab}
	10	115.2 ^{de}	210.0 ^c	71.88 ^c	69.8 ^c	1.56 ^{ab}	0.33 ^{ef}	0.58 ^{cd}	1.70 ^c
4.7	0	105.6 ^c	300.7 ^{cd}	81.23 ^{abc}	80.3 ^{abc}	1.17 ^{cd}	0.31 ^f	0.40 ^e	1.15 ^c
	5	145.6 ^b	450.5 ^a	84.49 ^{ab}	84.5 ^{ab}	1.36 ^{bc}	0.61 ^a	0.93 ^a	2.27 ^a
	10	128.0 ^{bcd}	269.0 ^d	77.98 ^{abc}	76.7 ^{bc}	1.58 ^a	0.55 ^b	0.70 ^b	1.90 ^{bc}
7.0	0	112.0 ^{de}	323.3 ^{bc}	81.65 ^{abc}	84.7 ^{ab}	1.36 ^{bc}	0.23 ^h	0.62 ^c	1.78 ^c
	5	105.6 ^c	349.0 ^{bc}	89.06 ^a	90.3 ^a	1.53 ^{ab}	0.25 ^{gh}	0.38 ^{ef}	1.27 ^{de}
	10	129.6 ^{bcd}	180.0 ^c	88.08 ^a	90.3 ^a	1.36 ^{bc}	0.29 ^{fg}	0.53 ^d	1.15 ^c

Means followed by the same superscript letters are not significantly different at P<0.05 according to Duncan's multiple test.

Data in the same Table (3) show that the interaction between irrigation with saline water and spraying plants with salicylic acid, had an impact on the water content of the leaves, expressed as RWC. The highest values were seen at treatment irrigated with 7.0 dSm⁻¹ and sprayed with 5.0 or 10.0 mM salicylic acid without differences between them. Relative water content of a leaf is a measurement the of 'water deficit' in the leaf, and may indicate a degree of water stress. Changes in RWC are proportional to changes in turgor pressure of the leaf, and so becomes an indirect measure of the change in turgor pressure under salinity treatments.

We can point out that plant height-RGR reached a maximum during the first year and there declined

during on the second year consistently as shown in Table (3). The decreasing value of height-RGR was due to greater contribution to leaf dry matter (LDW) or may be due to the spread of the roots, a similar finding was reported by Ismail, (1992) on *Calotropis procera*.

Nitrogen, phosphorus, potassium, sodium percentage and Total Chlorophyll in leaves (2nd years)

Table (4) indicates that both of N% and K% were higher with 10.0 mM than 0 or 5.0 mM SA under all levels of salinity. Phosphorus% data indicated that the most effective treatment was found when *Jatropha* was sprayed with 10.0 mM SA under irrigation with water of 4.7 dSm⁻¹. It is noted from Table (4) that sodium % in leaves under 5.0 mM SA treat-

ment was lower than that with 0 and 10 mM, under all levels of salinity. Abdi *et al.*, (2011) found that plants sprayed with SA showed lowest amount of Na accumulation. One of tolerance mechanisms may be exclusion of Na⁺ from the shoots by retaining it in root. Also, they suggested that alteration of mineral uptake from SA applications may be one of the mechanisms for alleviation of salt stress, and positively affected K⁺/Na⁺ ratio. Similar explanation was reported by Parida *et al.*, (2016). Also, War *et al.*, (2011) concluded that SA plays an important role in induction of plant defensive system

against a variety of biotic and abiotic stresses.

Concerning chlorophyll concentration, data indicated that the most effective treatment was found when *Jatropha* was irrigated with water of 4.7 dSm⁻¹, and sprayed with 5.0 mM SA. As stress increases, chlorophyll content decreases faster than other pigments (Bannari *et al.*, 2007). Jaleel *et al.*, (2009) mentioned that the decrease of leaf chlorophyll content under high salinity might be a reason to the destruction of pigments and the instability of the pigment-protein complex.

Table 4. Main and interactions effects of water salinity levels and SA concentrations on Nitrogen %, Phosphorus%, Potassium%, sodium % and Total Chlorophyll contents in leaves of *Jatropha* at early-growth stage (2nd years)

Treatments		N %	P %	K %	Na %	Total Chlorophyll mg g ⁻¹
EC (dSm ⁻¹)	SA (mM)					
Main effects of water salinity						
0.32		3.55 ^b	0.20 ^b	1.05 ^c	0.38 ^c	0.26 ^b
2.3		3.45 ^b	0.19 ^b	1.80 ^a	1.10 ^a	0.27 ^b
4.7		3.95 ^a	0.35 ^a	1.40 ^b	1.00 ^b	0.47 ^a
7.0		3.43 ^b	0.21 ^b	0.88 ^d	1.07 ^{ab}	0.19 ^c
Main effects of SA concentrations						
	0	3.14 ^c	0.25 ^b	0.87 ^b	0.95 ^b	0.20 ^c
	5	3.61 ^b	0.19 ^c	0.87 ^b	0.66 ^c	0.43 ^a
	10	4.04 ^a	0.28 ^a	2.07 ^a	1.06 ^a	0.26 ^b
Interaction effects of water salinity and SA concentrations						
0.32	0	3.36 ^b	0.29 ^c	0.48 ^c	0.44 ^f	0.23 ^{dc}
	5	3.30 ^c	0.12 ^g	0.52 ^e	0.28 ^g	0.26 ^{cd}
	10	4.00 ^a	0.20 ^d	2.16 ^a	0.42 ^f	0.30 ^{bc}
2.3	0	2.80 ^d	0.15 ^{fg}	1.14 ^d	1.00 ^{bc}	0.19 ^e
	5	3.36 ^{bc}	0.28 ^c	1.84 ^b	0.89 ^{cd}	0.29 ^{bc}
	10	4.20 ^a	0.15 ^{fg}	2.30 ^a	1.41 ^a	0.33 ^b
4.7	0	3.72 ^{abc}	0.33 ^b	1.22 ^d	0.98 ^{bc}	0.21 ^e
	5	3.92 ^a	0.19 ^{de}	0.62 ^e	0.76 ^{de}	0.89 ^a
	10	4.21 ^a	0.56 ^a	2.31 ^a	1.31 ^a	0.31 ^{bc}
7.0	0	2.67 ^d	0.21 ^d	0.64 ^e	1.40 ^a	0.18 ^e
	5	3.86 ^{ab}	0.16 ^{ef}	0.50 ^e	0.72 ^e	0.27 ^{cd}
	10	3.75 ^{abc}	0.26 ^c	1.50 ^c	1.10 ^b	0.11 ^f

Means followed by the same superscript letters are not significantly different at P<0.05 according to Duncan's multiple test.

Vegetative growth; seed yield and oil content (%) on early mature stage (4th year)

Effects of interaction between SA application and water salinity on some vegetative characters, TLA; RWC; RGR; LDW and seeds yield and oil content at early mature stage are listed in Table (5). Data shows that, the studied vegetative growth characters were significantly affected by interaction between SA concentrations and water salinity. The most effective treatments for growth characters were occurred when *Jatropha* shrubs were irrigated by saline water under level of 0.32 dSm⁻¹ (fresh water) but without significant difference with that irrigated with 4.7 dSm⁻¹ particularly, at 5.0 mM SA. It's clear that, SA application at 5.0 mM with either 0.32 or 4.7 dSm⁻¹ level increased fruit yield. These results are in agreement with those obtained on *Jatropha* by Hussein *et al.*, 2013. Application of 5 mM SA affected positively the growth characters at diluted sea water (4.7 dSm⁻¹) compared with

0 or 10.0 mM SA. The positive effect of SA on growth parameters could be attributed to its bio-regulator effect on physiological and biochemical processes in plants (EL Tayeb, 2005). Values of RWC as a character used to assess water status in leaf tissues indicated that there was no water deficits between fresh irrigated and salt-stressed leaves, which indicate that application of SA may lead to reduce salinity stress

In relation to oil % of seeds, data indicated that, the most effective treatment was found when *Jatropha* was sprayed with 5.0 mM SA and irrigated with 0.32 or 4.7dSm⁻¹. In general, average oil recovery of 25% by weight from seeds of *Jatropha* plant cultivated in different places (Tewfik *et al.*, 2012 and Pandey *et al.*, 2012). The oil content of *Jatropha* seeds of the actual study was determined quantitatively to be 20% based on dry seed weight. The variation in oil quantity might be due to change in climatic agricultural conditions (Hawash *et al.*, 2008).

Table 5. Main and interactions effects of water salinity levels and SA concentrations on some vegetative growth, seed yield and oil content (%) of *Jatropha* shrubs at early mature stage (4th years).

EC dSm ⁻¹	SA mM	TLA cm ²	RWC _L %	Height RGR cm month ⁻¹	LDW, g plant ⁻¹	Seed yield, g plant ⁻¹	Oil content %
Main effects of water salinity							
0.32		33948 a	85.15a	1.60a	293.29a	84.8a	14.4a
4.7		23611b	86.72a	1.64a	201.46b	82.6a	15.06a
Main effects of SA concentrations							
	0	27274.5a	83.9a	1.93a	312.62a	65.4a	14.1b
	5	30200.0b	88.09a	1.75a	250.93b	102.35a	17.35a
	10	18764.0c	85.79a	1.18b	178.58c	83.35b	12.8b
Interaction effects of water salinity and SA concentrations							
0.32	0	57520 a	83.70 a	2.58 a	496.13 a	72 cd	13.7c
	5	19050 d	86.36 a	1.26 cd	149.70d	98.7 ab	17.6a
	10	25274 c	85.40 a	0.96 d	234.05c	83.7 bc	12.0c
4.7	0	17229 dc	84.17 a	1.29 c	129.11d	58.8 d	14.5bc
	5	41350 b	89.83 a	2.25b	352.17b	106 a	17.1ab
	10	12254 c	86.18 a	1.40 c	123.11d	83 bc	13.6c

Means followed by the same superscript letters are not significantly different at P<0.05 according to Duncan's multiple test.

Nitrogen, phosphorus, potassium, sodium percentage and Total Chlorophyll in leaves (4th year):

Data in Table (6) show that Nitrogen percentage with 10.0 mM SA treatments was higher than 0 or 5.0 mM SA under water salinity of 4.7 dSm⁻¹, but Potassium percentage with 10.0 SA mM treatments was higher than 0 or 5 mM SA under all levels of water salinity. Data of Phosphorus percentage indicated that, the

most effective treatment was found when *Jatropha* irrigated under water of 0.32 dSm⁻¹, without addition of SA, while Sodium percentage with 5.0 mM SA treatment was lower than that with 0 and 10.0 mM, under 4.7 dSm⁻¹ water salinity. In relation to total chlorophyll, data indicate that, the most effective treatment was found when *Jatropha* was sprayed with 5.0 mM SA, under low levels of water salinity.

Table 6. Main and interactions effects of water salinity levels and SA concentrations on Nitrogen%, Phosphorus%, Potassium%, sodium % and Total Chlorophyll content on leaves of *Jatropha* at early-mature stage (4th year)

EC, dSm ⁻¹	SA mM	N %	P %	K %	Na %	Total Chlorophyll mg g ⁻¹
Main effects of water salinity						
0.32		3.10 ^a	0.27 ^a	0.72 ^a	0.46 ^a	0.32 ^a
4.7		3.35 ^a	0.07 ^b	0.46 ^b	0.97 ^a	0.35 ^a
Main effects of SA concentrations						
	0	2.99 ^b	0.16 ^a	0.52 ^b	0.6 ^b	0.29 ^b
	5	3.31 ^{ab}	0.1 ^b	0.38 ^c	0.7 ^b	0.38 ^a
	10	3.39 ^a	0.09 ^b	0.87 ^a	0.8 ^a	0.35 ^a
Interaction effects of water salinity and SA concentrations						
0.32	0	2.97 ^b	0.278 ^a	0.50 ^c	0.40 ^d	0.30 ^b
	5	3.25 ^{ab}	0.08 ^c	0.66 ^b	0.52 ^c	0.37 ^a
	10	3.08 ^b	0.126 ^b	1.00 ^a	0.46 ^{cd}	0.31 ^b
4.7	0	3.02 ^b	0.045 ^d	0.53 ^c	0.91 ^b	0.29 ^b
	5	3.36 ^{ab}	0.121 ^b	0.11 ^d	0.88 ^b	0.38 ^a
	10	3.69 ^a	0.07 ^c	0.74 ^b	1.14 ^a	0.39 ^a

Means followed by the same superscript letters are not significantly different at P<0.05 according to Duncan's multiple test.

In spite of higher *accumulation* of sodium in leaf tissue with increasing salinity (Tables 4 and 6), the RWC of leaf did not declined but it was increased due to high salinity. This artifact provide an indirect indication that Na⁺ is effectively compartmentalized in the vacuoles and this is required for osmotic adjustment, so it can be tacitly assumed that water relations and the ability to adjust the osmotic concentration play a vital role in overall growth and development of plants. In summary, results suggest that *Jatropha* can tolerate salinity by maintaining osmotic

balance and *ion* homeostasis. Same conclusion was recorded by Parida *et al.*, (2016).

Values of soil salinity

Soil salinity at the end of study for both studied stages: (the end of second year and immediately after fruits collecting), the EC in soil was measured and listed in Table (7). Data indicate that, the higher water salinity in irrigation water Leads to the higher residual soil salinity. So, each plant area was periodically leached dependent on the growth stage as described by (Hokam and Abo El-Soud, 2016).

Table 7. Values of soil salinity (Electrical conductivity) for early-growth and early mature stages

juvenile stage				
EC _{soil}	7.0 dSm ⁻¹	4.7 dSm ⁻¹	2.3 dSm ⁻¹	0.32 dSm ⁻¹
	7.64	5.11	2.61	0.96
Early mature stage				
EC _{soil}	--	4.7 dSm ⁻¹	--	0.32 dSm ⁻¹
		6.0		2.7

Conclusion

To avoid the competition with food production, marginal land and poor quality water must be targeted for producing bioenergy crops... SA application of 5 mM was positively affected vegetative growth characters, particularly at 4.7dSm⁻¹ salinity level. In mature stage, values of leaf relative water content character indicated that water deficits in leaf tissues did not exist between fresh irrigated and salt-stressed plants. The most potential vegetative growth and seed yield at mature stage were performed under treatment with fresh water compared to that of 7.2 or 4.7 dSm⁻¹. Application of 5.0 mM SA had a positive effect on both vegetative growth and capsule yield for 4.7dSm⁻¹ salinity treatment, compared to those for fresh water. It is recommended to interest with breeding for this shrub so can be overcome confusion and mystery resulting from genetic variation because original of plant still by seed.

References

Abdi, G.; Mohammadi, M. and Hedayat, M. (2011). Effect of salicylic acid on Na⁺ accumulation in shoot and roots of tomato in different K⁺ status. *J. Biol. Environ. Sci.*, 5(13): 31-35.

Allakhverdiev, S.I.; Sakamoto, A.; Nishiyama, Y.; Inaba, M. and Murata N. (2000). Ionic and Osmotic Effects of NaCl-Induced Inactiva-

tion of Photosystems I and II in *Synechococcus* sp. *Plant Physiology*. 123(3):1047-1056.

Antúnez, I. (2001). Relative growth rate in phylogenetically related deciduous and evergreen woody species. – *Oecologia* 128: 172–180.

Bannari, A. Khurshid, K. S.; Staenz, K. and Schwarz, J.W. (2007). A comparison of hyperspectral chlorophyll indices for wheat crop chlorophyll content estimation using laboratory reflectance measurements. *IEEE Trans. Geosci. Remote Sensing*, 45:3063-3074.

Bilal, S.; Mohammed-Dabo, I. A; Nuhu, M.; Kasim, S. A.; Almustapha I. H. and Yamusa Y. A. (2013). Production of biolubricant from *Jatropha curcas* seed oil. *Journal of Chemical Engineering and Materials Science*. 4: 72-79.

Crocker, A. (2013). *Renewable Energy in Egypt: Hydro, Solar and Wind*. Norton Rose Fulbright. <http://www.nortonrosefulbright.com/knowledge/publications/74735/renewable-energy-in-egypt>.

Dagar, J.C.; Tomar, O.S.; Kumar, Y. Bhagwan, H. Yadav, R.K. and Tyagi, N.K. (2006). Performance of some under-explored crops under saline irrigation in semiarid climate in Northwest India. *Land Degradation & Development* 17: 285-299.

El Tayeb, M. A. (2005). Response of barley grains to the interactive effect of salinity and salicylic acid. *Plant Growth Regu.*, 45: 215-224.

- Fujimaki, H. and Kikuchi N.N (2010). Drought and salinity tolerances of young *Jatropha*. *Int. J. Phys. Sci.* 24: 121-127.
- Hawash, S.; Kamal, N. and El Diwani, G. (2008). Egyptian *Jatropha* Oil Extraction for Biodiesel production. *AFINIDAD* 15: 470 – 474.
- Hoekstra, A.Y. and Chapagain. A.K. (2007). Water footprints of nations: water use by people as a function of their consumption pattern. *Water Resource Manage.* 21, 35-48.
- Hokam, E. M. and Abo El-soud, I. H. (2016). Evaluation of Growth Parameters for *Jatropha curcas* L. (Biofuel Plant) under Salinity and Water Stress Using Tensiometer for Irrigation Guiding. *Egyptian Journal of Soil Science*, 56: In press.
- Hussein, M.M.; El-Ashry, K. and El-Faham, S. Y. (2013). Effect of Salinity and MKP on Growth and Mineral Status of *Jatropha* Plants *Journal of Applied Sciences Research*, 9 (7): 4193-4203.
- Iqbal, M.; Ashraf, M. and Jamil, A. (2006). Seed enhancement with cytokinins: changes in growth and grain yield in salt stressed wheat plants. *Plant Growth Regul.* 50: 29-39.
- Ismail, A.M.A. (1992). Physiological Growth Analysis of the Life-History of *Calotropis procera* (Alt.) Alt. F. *Qatar Univ. Sci. J.* 12: 94- 100.
- Jaleel CA, Sankar B, Sridharan R, Panneerselvam R (2009). Soil salinity alters growth, chlorophyll content, and secondary metabolite accumulation in *Catharanthus roseus*. *Turk J Biol* 32(2): 79–83.
- Jewitt, G.P.W.; Wen, H.W.; Kunz, R.P. and Van Rooyen, A.M., (2009). Scoping Study on Water Use of Crops/Trees for Biofuels in South Africa. Report to the Water Research Commission, Wrc Report No. 1772/1/09 (ISBN 978-1-77005-884-2).
- Jones, Jr. J.B. (2001). *Laboratory Guide for Conducting Soil Tests and Plant Analysis*. CRC Press, Washington, DC., USA., ISBN-13: 9781420025293, p: 384.
- Jongschaap, R. E.; Corre, P. S. Bindra-ban, C. P. S. and Brandenburg, W.A. (2007). Claims and Facts on *Jatropha curcas* L. Report 158, Plant Research International, Wageningen.
- Khan, M.; Fatma. M.; Per, T.S.; Anjum, N.A. and Khan, N.A. (2015). Salicylic acid-induced abiotic stress tolerance and underlying mechanisms in plants. *Front Plant Sci.* 6: 462-468.
- Khan, M.R; Mehar, F.; Tasir S. P.; Anjum, N.A. and N. A. Khan (2015). Salicylic acid-induced abiotic stress tolerance and underlying mechanisms in plants. *Front Plant Sci.* 6: 462-469.
- Kotoky, R. A.; Rabha, A.; Gogoi, A.; Nath S. C. and Saikia, S. P. (2015). Correlation studies in seed traits, moisture and oil content and effect of hormones on flowering of *Jatropha curcas* L. *Brazilian Journal of Biological Sciences*, 2: 79-84.
- Licht, F.O. (2009). “2009 World Ethanol Production Growth to Hit Five-Year Low.” *World Ethanol & Biofuels Report.*”
- Maggio, A.; Reddy, M. P. and Joly R. J. (2000). Leaf gas exchange and solute accumulation in the halophyte *Salvadora persica* grown at moderate salinity. *Environ. Exp. Bot.* 44: 31–38.
- Mazumdar, B.C. and Majumder, K. (2003). *Methods of Physicochemical Analysis of Fruits*. Daya Publishing House Delhi, India.

- Najafian, S.; Khoshkhui M.; Tavallali V. and Saharkhiz M.J. (2009). Effect of salicylic acid and salinity in thyme (*Thymus Vulgaris L.*): Investigation on changes in gas exchange, water relations, and membrane stabilization and biomass accumulation. Aust. J. Basic. Applied Sci., 3: 2620-2626.
- Niu, G., D. Rodriguez, J. Mendoza, M. Jifon and G. Ganjegunte, (2012). Responses of *Jatropha curcas* to Salt and Drought Stresses. International Journal of Agronomy, Vol. 2012. Article ID 632026, 1 - 7.
- Pandey, V.C.; Singh, K.; Singh, J.S.; Kumar, A.; Singh, B. and Singh, R.P. (2012). *Jatropha curcas*: A Potential Biofuel Plant for Sustainable Environmental Development. Renewable and Sustainable Energy Reviews, 16: 2870-2883.
- Parida, A.K.; Veerabathini, S.K.; Kumari, A. and Agarwal, P.K. (2016). Physiological, Anatomical and Metabolic Implications of Salt Tolerance in the Halophyte *Salvadora persica* under Hydroponic Culture Condition. Front Plant Sci. Article 7: 1-18.
- Pérez-Harguindeguy N., Díaz S., Garnier E., *et al.* (2013). New handbook for standardised measurement of plant functional traits worldwide. Australian Journal of Botany, 61, 167-234.
- Saleh, B. (2012). Salt stress alters physiological indicators in cotton (*Gossypium hirsutum L.*). Soil Environ. 31(2): 113-118.
- Shibghatallah1, M. A. H.; Khotimah, S.N.; Suhandono, S.; Viridi1,S., and Kesuma,T.(2013). Measuring Leaf Chlorophyll Concentration from Its Color: A Way in Monitoring Environment Change to Plantations”, physics. bio-ph 2: 1305.1148.
- Shraiy, A.M.E. and Hegazi, A.M. (2009). Effect of acetylsalicylic acid, indole-3-butyric acid and gibberellic acid on plant growth and yield of pea (*Pisum Sativum L.*). Aust. J. Basic Appl. Sci., 3: 3514–3523.
- Tewfik, S.R.; Hawash, S.I.; Atteya, N. K.; Diwani G. I. and Farag, I.H. (2012). Techno-economic Appraisal of Biodiesel from *Jatropha curcas*: An Egyptian Case Study. Journal of Agricultural Science and Technology 2: 287-297.
- Villar-Salvador, P.; Puértolas, J.; Cuesta, B.; Peñuelas, J.L.; Uscola, M.; Heredia-Guerrero, N. and Rey Benayas, J.M. (2012). Increase in size and nitrogen concentration enhances seedling survival in Mediterranean plantations. Insights from an ecophysiological conceptual model of plant survival. *New For* 43:755 –770.
- War, A.R; Paulraj, M.G.; War, M.Y. and Ignacimuthu S. (2011). Role of salicylic acid in induction of plant defense system in chickpea (*Cicer arietinum L.*) Plant Signaling & Behavior 6:11, 1787-1792.

تأثير الاضافة الخارجية لحمض السالسيك علي الخصائص المورفولوجية والكيميائية لنبات الجتروفا المروي بماء مالح

إسلام حسن أبوالسعود^١ وعصام محمد حكام^٢

^١ قسم البساتين، كلية الزراعة، جامعة قناة السويس

^٢ قسم الأراضي والمياه، كلية الزراعة، جامعة قناة السويس

الملخص

الملوحة واحده من اهم العوامل المؤثره علي انتاجية المحاصيل، اجريت دراسة حقلية علي مرحلتي نمو للنبات الجتروفا (نبات وقود حيوي) ، مرحلة النمو المبكرة (السنة الاولى الي السنة ثانية) و مرحلة الاثمار المبكرة (السنة الثالثة الي السنة الرابعة)، تم تحضير ثلاث مستويات من الماء المالح بتخفيف ماء البحر (تم تجميعه من قناة السويس) وهي ٢,٣ و ٤,٧ و ٧ dSm⁻¹. بالإضافة الي الماء العذب ، تم رش نباتات الجتروفا بثلاث تركيزات من حمض السالسيك (١٠-٥-٠٠ - مليمول) ، النتائج المتحصل عليها في مرحلة النمو المبكرة اشارت الي ان الري بالماء العذب اعطي افضل نمو لصفات النمو الخضريه (المساحة الكلية للاوراق ، محتوى الرطوبة النسبية للاوراق ، معدل النمو النسبي والوزن الجاف للاوراق) ولكن بدون فروق معنوية مع ماء ري عند تركيز ٢,٣ و ٤,٧ dSm⁻¹. الرش بحمض السالسيك عند تركيز ٥ مليمول زاد من قياسات النمو الخضريه في وجود ماء ري بتركيز ملوحة ٤,٧ dsm⁻¹. وتفاوتت هذه علي اضافة ٠ او ١٠ مليمول حمض سالسيك.

فيما يخص النباتات في مرحلة الاثمار المبكرة كان لاضافة ٥ مليمول من حمض السالسيك تأثير ايجابي علي النمو الخضري والثمري تحت معاملة ماء الري ٤,٧ dSm⁻¹. مقارنة بالكنترول ، قراءات محتوى الرطوبة النسبي للاوراق اشارت الي عدم وجود فروق في رطوبة الاوراق للنباتات المزروعة في الماء العذب او الماء المالح مما يدل علي ان الرش بحمض السالسيك قد ادي الي تجنب الاجهء الملحي، الرش بحمض السالسيك ادي الي زيادة تركيز الكلورفيل، بصفة عامة رش شجيرات الجتروفا بحمض السالسيك كان له تأثير ايجابي في تقليل التأثير السلبي للماء المالح.