

## **Effect of GA<sub>3</sub>, Urea and Yeast Spraying on Fruiting of Flame Seedless Grapevines under Sandy Soil Conditions**

**El-Halaby, E.H.S.<sup>1</sup>; \*A.M. El-Salhy<sup>1</sup>; M.M. Al-Wasfy<sup>2</sup> and R.A. Ibrahim<sup>1</sup>**

<sup>1</sup>Pomology Dept., Fac. Agric., Assiut University, Assiut.

<sup>2</sup>Hort. Dept., Fac. Agric., South Valley University, Qena.

\*E-mail: [abdelfattah.elsalhy@agr.au.edu.eg](mailto:abdelfattah.elsalhy@agr.au.edu.eg)

---

### **Abstract:**

This investigation was carried out during three seasons i.e. 2010, 2011 and 2012 on Flame Seedless grapevines grown in sandy soil in El-Karnak vineyard for table grapes production, Luxor governorate, Egypt. Ten combined treatments of GA<sub>3</sub>, urea and active dry yeast spraying at various stage of berry development were evaluated. The experimental vines were arranged in a complete randomized design with three replications per treatment, two vines each. From the results of this investigation it could be concluded that spraying with GA<sub>3</sub> seven times, once at pre-bloom (5 ppm), thrice at full-bloom (2.5, 5 and 7.5 ppm) and other thrice when the berry at (6 mm) pea stage (30, 30 and 20 ppm). As well as, combined spraying GA<sub>3</sub> four times once at pre-bloom and thrice at full-bloom plus 0.4% active dry yeast when the berry at pea stage to obtain heavy and less compact cluster and hasten the ripening with fairly good Flame Seedless berries quality. In addition, it could be used urea and yeast instead of GA<sub>3</sub> in grape production to overcome the adverse GA<sub>3</sub> effects especially for colored cultivars.

---

**Keywords:** Growth regulators, GA<sub>3</sub>, urea, active dry yeast, grapevines, thinning.

---

**Received on:** 11/3/2015

**Accepted for publication on:** 21/3/2015

**Referees:** Prof. Ahmed M. El-sese

Prof. Ahmed H. Abdel-Aal

### **Introduction:**

Grapes (*Vitis vinifera* L.) are considered the first major fruit crop in its production all over the world, for being of an excellent favour, nice taste and high nutritional value. In Egypt grapes rank second among fruit crops while citrus being the first. The total planted area attained about 188543 feddans with an average of 1378815 tons (Annual Statistical of the Ministry of Agriculture, 2012).

The merit of enhancing ripening and improving quality of early grapevine cultivars are clear for increasing the opportunity of exportation by using plant growth substances. The improving early grapes is very important either for local consumption markets or exportation to external markets. Plant growth substances play a major role in plant growth and development. GA<sub>3</sub> still used to increase cluster length, berry size and for thinning bunch berries in Seedless grape cultivars (Orth, 1990; Colapetra *et al.*, 1995; El-Hammady *et al.*, 1998; Marzouk and Kassem, 2002; Williams and Ayars, 2005; Selim, 2007 and Zoffoli *et al.*, 2009).

Berry thinning has been used to obtain a good cluster with highest berry weight and fastest ripening. Bunch thinning is done as a regular cultural treatment or spray of chemicals at pre-bloom, peak bloom and fruit set stages. The practice is done to reduce cluster compactness and to improve the productivity and berry quality. The thinning necessary depended on the cultivar as well as sunshine, temperature and nutrient supply (Dhillon *et al.*, 1992; Poni, 2003 and Ahmed *et al.*, 2004).

Recently urea spraying at pre-bloom or full bloom has been used to

reduce the berry set percentage and consequently to induce berry thinning (Ahmed *et al.*, 2004; El-Salhy *et al.*, 2009 and Fawzi *et al.*, 2014). The bio-fertilizer active dry yeast was enhanced grape yield and berry quality where, yeast contains some natural growth regulators, some important nutrients and some common amino acids (Moor, 1979; Idso *et al.*, 1995; El-Salhy *et al.*, 2011 and Fawzi *et al.*, 2014).

The purpose of this study was to add more light on the effect of spraying GA<sub>3</sub>, urea and active dry yeast on yield, clusters attributes and berry quality of Flame Seedless grape cultivar under sandy soil.

### **Materials and Methods:**

The present work was conducted through three successive seasons of 2010, 2011 and 2012 on 60 uniform vigour seven years-old Flame Seedless grapevines. The vines were grown in El-Karnak vineyard for table grapes production-Luxor Governorate, Egypt. They grown in sandy soil at 2x3 meters apart under drip irrigation system with salty water 1500 ppm. All vines received the standard agricultural practices that are used in the vineyard including spraying dormex, soil fertilization, irrigation and pest control. The Spanish Barron system was used as a trellising system. The vines were cane pruned (68 eyes/vine were left, 10 canes x 6 buds/cane plus 4 renewal spurs with 2 buds). The pruning was done during the second week of January each season. Crop load at all vines was adjusted to 25 clusters/vine after berry set. The chosen vines were divided into ten different treatments including the control. The experimental vines were arranged in a complete

randomized block design with three replications per treatment two vines each. Thus, the treatments were as follow:

- 1- Control (sprayed with water only).
- 2- GA<sub>3</sub> at 5 ppm sprayed when cluster length was about (10-12 cm) for elongation.
- 3- GA<sub>3</sub> at 2.5 ppm plus 1% urea when cluster length was about (10-12 cm) for elongation.
- 4- 2% urea sprayed when cluster length was about (10-12 cm) for elongation.
- 5- GA<sub>3</sub> four times spraying at 5 ppm for elongation followed by thrice GA<sub>3</sub> at 2.5, 5 and 7.5 ppm sprayed during full bloom (80, 100 and 120% of the flowers caps, dropped) during the successive three days, respectively for berry thinning.
- 6- GA<sub>3</sub> three times spraying, once at 5 ppm for elongation followed by twice GA<sub>3</sub> at 2.5, 5 ppm and 1% urea for berry thinning.
- 7- GA<sub>3</sub> at 5 ppm for elongation followed by 2% urea for berry thinning.
- 8- GA<sub>3</sub> seven times spraying, once at 5 ppm for elongation, followed by thrice GA<sub>3</sub> at 2.5, 5, 7.5 ppm for thinning and other thrice 30, 30, 20 ppm of GA<sub>3</sub> when berry diameter reached about 6 mm (pea stage) for sizing.
- 9- GA<sub>3</sub> six times spraying, once at 5 ppm for elongation, followed by thrice 2.5, 5 & 7.5 ppm of GA<sub>3</sub> for berry thinning and then twice GA<sub>3</sub> at 15, 15 ppm and 0.2% yeast for sizing.
- 10- GA<sub>3</sub> four times spraying, once at 5 ppm for elongation followed by thrice 2.5, 5 & 7.5 ppm of

GA<sub>3</sub> for berry thinning and then 0.4% yeast for sizing.

The substances used in this experiment were GA<sub>3</sub>, urea and active dry yeast to study the impact of spraying them on yield, cluster attributes and berry quality. GA<sub>3</sub> (Gibberellic acid), and low biuret urea (46%) were prepared before spraying by dissolved the define amount in water based. Active dry yeast was prepared by dissolved the define amount in warm water (38°C) followed by addition of 0.3% Egyptian treacle (as source of sugar) and left for two hours before spraying for activating. All chemicals were sprayed at same date by using a hand sprayer to the runoff. The percentage of berry set was estimated by caging two cluster per vine in perforated white cheese bags after the first spraying. Such bags were removed for chemical spraying at blooming, the percentage of berry set was calculated as follow:

$$\text{Berryset \%} = \frac{\text{No. of berries/cluster}}{\text{No. of total flowers/cluster}} \times 100$$

At harvest time (when TSS of berry juice in the check treatment reached 13-14% brix), the clusters were harvested, weighed and yield/vine (kg) was recorded. Two clusters were taken at random from yield of each vine and the following characteristics were determined.

Cluster weight (g), cluster length (cm) and number of berries per cluster, then cluster compactness coefficient according to Winkler *et al.* (1974), as well as shot berries percentage.

In addition berry quality in terms of berry weight, TSS, total ti-

tratable acidity and reducing sugars % according to A.O.A.C. (1985), as well as total anthocyanin according to Markham (1982).

All obtained data were tabulated and statistically analysed according to Gomez and Gomez (1984) and Snedecor and Cochran (1990) using the L.S.D. test for distinguishing the significance differences between various treatment means.

## **Results:**

### **1- Berry set percentage and yield:**

Data presented in Table (1) shows the effect of spraying with GA<sub>3</sub>, low biuret urea and active dry yeast on berry set percentage, shot berries percentage and yield of Flame Seedless grapevines in 2010, 2011 and 2012 seasons. The results took similar trend during the three studied seasons. It is obvious from the obtained data that the single or combined spraying of GA<sub>3</sub> and urea at full bloom significantly decreased the berry set percentage and shot berries percentage compared to untreated vine. The decrement percentage of berry set and shot berries percentage due to spray of GA<sub>3</sub> or urea single or combination compared to unsprayed one were attained 23.36, 25.48 & 25.02% and 76.11, 79.75 & 82.16% as an average of the three studied seasons, respectively. On other hand, single or combined GA<sub>3</sub> and active dry yeast spraying after berry set significantly increased the yield/vine compared to other treatments and unsprayed ones (control). GA<sub>3</sub> spraying suppressed the yeast spraying concerning yield/vine, but had insignificant differences compared to active dry yeast. The increment percentage of the yield was attained 27.33, 21.71

& 22.99% as an average of the three studied seasons, due to GA<sub>3</sub> (T<sub>8</sub>), GA<sub>3</sub> plus active (T<sub>9</sub>) and active dry yeast after berry set (T<sub>10</sub>) follow GA<sub>3</sub> for elongation and thinning compared to untreated one (control), respectively.

### **2- Cluster characteristics:**

The effect of tested treatments on cluster characteristics during the three studied seasons are shown in Tables (2 and 3). It is evident that all treatments improved the cluster traits. Using GA<sub>3</sub> or urea at pre-bloom significantly increased the cluster length, whereas, using them at full-bloom significantly decreased the berries number per cluster compared to untreated one (control), hence significantly decreased compactness coefficient of cluster and produced loose clusters. The increment percentage of cluster length was attained (16.09, 12.84 and 11.24% as an average of the three studied seasons) due to spray GA<sub>3</sub>, GA<sub>3</sub> plus urea and urea at pre-blooming compared to untreated ones, respectively. On the other hand, the decrement percentage of berries number per cluster was attained 19.97, 22.74 and 18.47% as an average of the three studied seasons, due to spray GA<sub>3</sub>, GA<sub>3</sub> plus urea and urea at full-blooming followed GA<sub>3</sub> spraying for elongation (T<sub>2</sub>), respectively. Hence, the corresponding decrement percentage of cluster compactness coefficient was 28.66, 31.10 and 27.32% as an average of the three studied seasons, respectively.

Moreover, spraying either GA<sub>3</sub> or active dry yeast after berry set singly or combination were increased the cluster weight compared to control or other treatments. Insignificant

differences were observed between used GA<sub>3</sub> or urea for cluster elongation and berry thinning, as well as used GA<sub>3</sub> or yeast after berry set for sizing. The increment percentage of cluster weight was attained 27.48, 21.98, 23.61% as an average of the three studied seasons, due to GA<sub>3</sub>, plus active dry yeast and yeast spraying after berry set, following GA<sub>3</sub> (T<sub>2</sub>) and GA<sub>3</sub> spraying at full blooming for berry thinning (T<sub>5</sub>), compared to untreated one (control), respectively. Therefore, it can be concluded that single or combined spraying of GA<sub>3</sub> or urea at berry blooming and full blooming, then following single or combined spraying of GA<sub>3</sub> or active dry yeast after berry set was the best tool to produce loose clusters with heavy weight.

### **3- Berry quality:**

Data of various berry characteristics as affected by different studied treatments during 2010, 2011 and 2012 seasons are presented in Tables (3 and 4). The data indicated that GA<sub>3</sub> and urea spraying at pre-bloom and full bloom, and followed by GA<sub>3</sub> or active dry yeast spray after berry set significantly improved the Flame Seedless grapes quality in terms of berry weight, total soluble solids, re-

ducing sugars and anthocyanin in berry skin compared to untreated ones. The increment percentage of berry weight was 57.87, 54.33 and 52.32% as an average of the three studied seasons, due to GA<sub>3</sub> (T<sub>8</sub>), GA<sub>3</sub> plus active dry yeast (T<sub>9</sub>) or active dry yeast sprayed after berry set (T<sub>10</sub>) followed by GA<sub>3</sub> for elongation and berry thinning compared to untreated one (control), respectively. As well as, the corresponding increment percentage of total soluble solids was 2.31, 5.93 & 7.24%, respectively. Moreover, the increment percentage of anthocyanin in berry skin was 18.39 & 28.74% as an average of the three studied seasons, due to either GA<sub>3</sub> plus active dry yeast (T<sub>9</sub>) or yeast (T<sub>10</sub>) compared to GA<sub>3</sub> spraying (T<sub>8</sub>), respectively. Using singly yeast spraying or GA<sub>3</sub> plus yeast for sizing had the highest berry weight and size with best chemical juice quality compared to GA<sub>3</sub> only. Therefore, it can be concluded that could be used active dry yeast as a tool for grape berry sizing instead of GA<sub>3</sub> to overcome the adverse effects due to GA<sub>3</sub> in grape production, i.e. delaying the berry ripening and reducing full coloration, especially with colored cultivars.

**Table (1): Effect of GA<sub>3</sub>, urea and yeast spraying on berry set %, shot berries % and yield of Flame Seedless grapevines during 2010, 2011 and 2012 seasons.**

No.	Treat.	Ch.	Berry set %				Shot berries %				Yield kg/vine			
			2010	2011	2012	Mean	2010	2011	2012	Mean	2010	2011	2012	Mean
1	Control (water)		14.95	14.61	15.78	15.11	12.70	12.08	13.90	12.89	7.9	7.2	8.4	7.83
2	5 ppm GA <sub>3</sub>		15.10	14.58	15.52	15.07	3.45	3.52	2.98	3.32	8.5	8.0	9.2	8.57
3	2.5 ppm GA <sub>3</sub> + 1% urea		15.31	14.86	15.49	15.22	3.40	3.28	3.17	3.22	8.6	7.9	9.3	8.60
4	2% urea		14.58	14.23	14.84	14.55	3.71	3.62	3.88	3.56	8.0	7.7	8.8	8.17
5	T <sub>2</sub> + 2.5, 5, 7.5 ppm GA <sub>3</sub>		11.53	11.28	11.93	11.58	2.80	3.00	3.45	3.08	8.6	8.1	9.4	8.70
6	T <sub>2</sub> + 2.5, 5 ppm GA <sub>3</sub> + 1% urea		11.24	11.05	11.50	11.26	2.80	2.65	2.40	2.61	8.5	8.1	9.3	8.63
7	T <sub>2</sub> + 2% urea		11.20	11.35	11.45	11.33	2.40	2.35	2.16	2.30	7.9	7.7	8.9	8.17
8	T <sub>5</sub> + 30, 30, 20 ppm GA <sub>3</sub>		11.78	11.30	11.83	11.64	2.50	2.30	2.36	2.39	9.7	9.3	10.9	9.97
9	T <sub>5</sub> + 15, 15 ppm GA <sub>3</sub> + 0.2% yeast		11.36	11.41	11.66	11.48	2.80	2.35	3.23	2.79	9.3	8.9	10.4	9.53
10	T <sub>5</sub> + 0.4% yeast		11.57	11.33	11.60	11.50	2.23	2.15	3.30	2.56	9.4	9.0	10.5	9.63
L.S.D. at 5%			0.75	0.70	0.83		0.22	0.28	0.18		0.44	0.48	0.61	

T<sub>2</sub> (5 ppm GA<sub>3</sub> for cluster elongation)

T<sub>5</sub> (T<sub>2</sub> plus 2.5, 5, 7.5 ppm GA<sub>3</sub> for thinning)

T<sub>8</sub> (T<sub>5</sub> plus 30, 30, 20 ppm GA<sub>3</sub> for sizing)

**Table (2): Effect of GA<sub>3</sub>, urea and yeast spraying on cluster weight, No. of berries/cluster and cluster length of Flame Seedless grapevines during 2010, 2011 and 2012 seasons.**

No.	Treat.	Ch.	Cluster weight (g)				No. berries/cluster				Cluster length (cm)			
			2010	2011	2012	Mean	2010	2011	2012	Mean	2010	2011	2012	Mean
1	Control (water)		314.3	288.5	335.0	312.6	136.2	131.7	133.3	133.7	16.0	16.32	16.51	16.28
2	5 ppm GA <sub>3</sub>		338.6	317.7	366.8	341.0	137.1	133.8	134.3	135.1	18.8	18.91	19.00	18.90
3	2.5 ppm GA <sub>3</sub> + 1% urea		342.5	314.8	370.6	342.6	138.2	134.6	135.8	136.2	18.40	18.32	18.38	18.37
4	2% urea		322.0	305.9	342.3	323.4	132.6	130.3	128.2	130.4	18.03	18.15	18.16	18.11
5	T <sub>2</sub> + 2.5, 5, 7.5 ppm GA <sub>3</sub>		343.4	324.5	376.3	348.1	107.2	108.1	105.8	107.0	18.60	18.18	18.14	18.31
6	T <sub>2</sub> + 2.5, 5 ppm GA <sub>3</sub> + 1% urea		337.8	323.0	369.5	343.4	104.2	102.2	103.5	103.3	18.33	18.25	18.40	18.33
7	T <sub>2</sub> + 2% urea		317.0	308.6	357.8	327.8	109.0	108.3	109.8	109.0	18.07	18.06	18.70	18.28
8	T <sub>5</sub> + 30, 30, 20 ppm GA <sub>3</sub>		385.3	373.9	436.3	398.5	108.0	105.6	108.6	107.5	18.50	18.08	18.65	18.41
9	T <sub>5</sub> + 15, 15 ppm GA <sub>3</sub> + 0.2% yeast		371.0	354.5	418.4	381.3	105.0	102.5	105.8	104.4	18.27	18.10	18.25	18.21
10	T <sub>5</sub> + 0.4% yeast		377.4	361.5	420.2	386.4	108.8	106.4	110.0	108.4	18.20	18.15	18.30	18.22
L.S.D. at 5%			16.69	15.87	19.88		5.47	6.18	5.88		0.73	0.85	0.89	

T<sub>2</sub> (5 ppm GA<sub>3</sub> for cluster elongation)

T<sub>5</sub> (T<sub>2</sub> plus 2.5, 5, 7.5 ppm GA<sub>3</sub> for thinning)

T<sub>8</sub> (T<sub>5</sub> plus 30, 30, 20 ppm GA<sub>3</sub> for sizing)

**Table (3): Effect of GA<sub>3</sub>, urea and yeast spraying on cluster compactness coefficient, 25 berries weight and TSS of Flame Seedless grapes during 2010, 2011 and 2012 seasons.**

No.	Treat.	Ch.	Compactness coefficient				25 Berries weight (g)				TSS %			
			2010	2011	2012	Mean	2010	2011	2012	Mean	2010	2011	2012	Mean
1	Control (water)		8.45	8.69	8.06	8.20	50.23	49.85	55.58	51.89	13.80	14.00	13.67	13.82
2	5 ppm GA <sub>3</sub>		7.29	7.10	7.05	7.15	54.43	53.40	62.63	56.82	14.67	14.80	14.48	14.65
3	2.5 ppm GA <sub>3</sub> + 1% urea		7.47	7.37	7.40	7.41	55.77	52.85	64.51	57.71	14.67	14.60	14.50	14.54
4	2% urea		7.38	7.16	7.08	7.21	51.82	50.70	60.03	54.18	15.00	14.80	14.80	14.87
5	T <sub>2</sub> + 2.5, 5, 7.5 ppm GA <sub>3</sub>		5.75	5.94	5.85	5.85	71.73	71.05	81.72	74.83	14.87	14.67	14.67	14.74
6	T <sub>2</sub> + 2.5, 5 ppm GA <sub>3</sub> + 1% urea		5.69	5.59	5.66	5.65	71.83	71.25	81.81	74.96	15.13	14.80	14.80	14.91
7	T <sub>2</sub> + 2% urea		6.03	5.98	5.88	5.96	65.32	64.60	73.95	67.96	14.80	14.82	14.90	14.84
8	T <sub>5</sub> + 30, 30, 20 ppm GA <sub>3</sub>		5.88	5.81	5.85	5.85	79.52	77.38	88.86	81.92	14.25	14.18	14.00	14.14
9	T <sub>5</sub> + 15, 15 ppm GA <sub>3</sub> + 0.2% yeast		5.77	5.64	5.81	5.74	77.40	76.20	86.64	80.08	14.60	14.60	14.73	14.64
10	T <sub>5</sub> + 0.4% yeast		5.99	5.88	5.85	5.91	76.65	75.10	85.38	79.04	14.87	14.80	14.80	14.82
L.S.D. at 5%			0.33	0.38	0.41		2.93	2.39	3.56		0.58	0.39	0.45	

T<sub>2</sub> (5 ppm GA<sub>3</sub> for cluster elongation)

T<sub>5</sub> (T<sub>2</sub> plus 2.5, 5, 7.5 ppm GA<sub>3</sub> for thinning)

T<sub>8</sub> (T<sub>5</sub> plus 30, 30, 20 ppm GA<sub>3</sub> for sizing)

**Table (4): Effect of GA<sub>3</sub>, urea and yeast spraying on some chemical constituents of Flame Seedless grape berries during 2010, 2011 and 2012 seasons.**

No.	Treat.	Ch.	Titratable acidity %				Reducing sugars %				Anthocyanin mg/g			
			2010	2011	2012	Mean	2010	2011	2012	Mean	2010	2011	2012	Mean
1	Control (water)		0.49	0.52	0.50	0.50	10.64	10.93	10.73	10.77	0.90	0.92	0.98	0.93
2	5 ppm GA <sub>3</sub>		0.50	0.51	0.50	0.50	11.34	11.60	11.62	11.52	0.94	0.96	1.02	0.97
3	2.5 ppm GA <sub>3</sub> + 1% urea		0.46	0.48	0.47	0.47	11.47	11.75	11.50	11.57	0.95	0.97	1.04	0.99
4	2% urea		0.46	0.47	0.46	0.46	11.68	11.83	11.76	11.76	0.99	1.02	1.08	1.03
5	T <sub>2</sub> + 2.5, 5, 7.5 ppm GA <sub>3</sub>		0.46	0.48	0.47	0.47	11.65	11.95	11.75	11.78	0.96	0.96	1.02	0.98
6	T <sub>2</sub> + 2.5, 5 ppm GA <sub>3</sub> + 1% urea		0.45	0.46	0.46	0.46	11.93	12.11	11.70	11.91	1.02	1.05	1.11	1.06
7	T <sub>2</sub> + 2% urea		0.45	0.45	0.46	0.45	11.59	11.93	11.82	11.78	1.08	1.11	1.16	1.12
8	T <sub>5</sub> + 30, 30, 20 ppm GA <sub>3</sub>		0.52	0.55	0.53	0.53	11.41	11.76	11.38	11.52	0.85	0.86	0.90	0.87
9	T <sub>5</sub> + 15, 15 ppm GA <sub>3</sub> + 0.2% yeast		0.48	0.50	0.48	0.49	11.36	11.60	11.58	11.51	0.94	1.07	1.08	1.03
10	T <sub>5</sub> + 0.4% yeast		0.45	0.46	0.45	0.45	11.85	11.80	11.75	11.73	1.08	1.11	1.16	1.12
L.S.D. at 5%			0.02	0.03	0.02		0.51	0.55	0.48		0.05	0.04	0.05	

T<sub>2</sub> (5 ppm GA<sub>3</sub> for cluster elongation)

T<sub>5</sub> (T<sub>2</sub> plus 2.5, 5, 7.5 ppm GA<sub>3</sub> for thinning)

T<sub>8</sub> (T<sub>5</sub> plus 30, 30, 20 ppm GA<sub>3</sub> for sizing)

### Discussion and Conclusion:

GA<sub>3</sub> has been routinely used for Seedless grape production to increase berry and cluster weight, and cause thinning of clusters. The effect of GA<sub>3</sub> depends on date of treatments and concentration applied (Perez *et al.*, 2000 and Casanova *et al.*, 2009).

GA<sub>3</sub> spraying at full bloom decreased berry set since its role in flower dropping, causing reduction of berry set induce a reduction of berries number of cluster (Lu *et al.*, 1995; Dokoozlian and Peacock, 2001; Selim, 2007; El-Salhy *et al.*, 2009 and Abu-Zahra, 2010). The positive action of GA<sub>3</sub> on stimulating cell elongation process,

enhancing the water absorption and stimulating the biosynthesis of proteins which will lead to increase the cluster length, as well as, berry size and weight. GA<sub>3</sub> spraying three times was more effective in improving the cluster traits, since decreasing the berries number, whereas increasing length and weight of cluster consequently significantly decreased the compactness coefficient (Roper and Williams, 1989; Lu *et al.*, 1995; El-Salhy *et al.*, 2009 and Abu-Zahra, 2010).

The results are on line with those obtained by the investigators, Ezzahouani *et al.* (1985), Orth (1990), Lu *et al.* (1995), Dokoozlian and Peacock (2001), Selim (2007), El-Salhy *et al.* (2009) and Abu-Zahra (2010). They revealed that there is a potential benefit from GA<sub>3</sub> treated grapevine in the commercial production of Seedless grapes for its effective influence on yield and quality.

In addition, the positive action of urea as nitrogen source and producing new tissues that water and nutrients absorption induce more vegetative growth that shifted the balance of competition between reproductive growth and vegetative organs in favor of the latter. Nitrogen has many function in all division, the synthesis of proteins, protoplasm, enzymes and organic compounds as nucleoproteins, amino acid and chlorophyll (Nijjar, 1985). Low buiret urea differed significantly from control in term of fruit set and fruit thinning percentage. The reasons may be the interference with fertilization of the ovary of phytotoxicity in the peduncle region (Byers and Lyons, 1985; Guirguis *et al.*, 1996 and Ahmed *et al.*, 2004). There was a remarkable improving on berry quality expressed on increasing the berry weight, total soluble solids,

reducing sugars and anthocyanin contents as berry thinning, such findings might be due to induce a decreasing of berries number which increasing the available amount of organic foods required for each individual berry hence induce advance the berry ripening. The results of urea on improving yield and berry quality of grapevines was supported by many authors such as El-Moursy *et al.* (1993), Abdel-Hady (1995), Ahmed *et al.* (2004), El-Salhy *et al.* (2009) and Fawzi *et al.* (2014).

Recently, active dry yeast was found to enhance grape yield and berry quality. Yeast has high content of mineral particularly, N, P and K, proteins, vitamin B and natural hormone, i.e. cytokinin and IAA. The improving effect of yeast application was attributed to axuins, hormones, vitamins, chelating agents and enzyme produced, which have stimulatory effects on cell division and enlargement, nutrient uptake, protein synthesis and improves net photosynthese (Moor, 1979 and Idso *et al.*, 1995). These effects induce an improving of hormones and accumulation of carbohydrates consequently raising sugars and anthocyanin contents in berry induce advancing of the berry ripening. It known that the earliest production are the most important target for export and marketing. The results dealing with the effect of yeast spraying on grapevine fruiting are in harmony with those of Hassan (2002), Omran and Abdel-Latif (2003), El-Akkad (2004), Omran *et al.* (2005), El-Salhy *et al.* (2011) and Fawzi *et al.* (2014). They concluded that spraying yeast significantly improved of berry quality in terms of increasing the berry size, TSS, reducing sugars and anthocyanin contents and de-

creasing the total acidity then induce advance the berry ripening.

On the light of the previous results, it could be recommended that spraying of GA<sub>3</sub> four times, once at pre-bloom (5 ppm), thrice at full-bloom (2.5, 5 & 7.5 ppm) plus 0.4% active dry yeast when the berry diameters about 6 mm (pea stage), as well as GA<sub>3</sub> seven times, once at pre-bloom, thrice at full-bloom and thrice (30, 30 & 20 ppm) when the berry of pea stage. In addition, can be used 2% low biuret urea at pre-bloom and full bloom, as well as 0.4 active dry yeast when the berry at pea stage. Using urea and yeast more effective to overcome the adverse effective of using GA<sub>3</sub> at high concentration, i.e. delay the berry ripening and reduce full coloration. These treatments very necessary to produce heavy and less compact cluster and hasten the ripening as well as improving the weight, size, colouration and taste of Flame Seedless berries. These advantage will eventually enable growers to obtain highly marketable surrounding and overseas markets.

#### References:

- Abdel-Hady, A.M. (1995): Response of Roomy Red grapevines to foliar sprays of urea and boron. Ph.D. thesis, Fac. of Agric., Minia Univ., Egypt.
- Abu-Zahra, T.R. (2010): Berry size of Thompson Seedless as influenced by the application of gibberellic acid and cane girdling. Pak. J. Bot., 42 (3): 1755-1760.
- Ahmed, W.; M. Junaid; S. Amin and M. Nafees (2004): Low biuret urea application at different phenophases of bunch to improve productivity of perlette grapes inter. J. of Agri. & Bio. 6 (2): 418-419.
- Annual Reports and Agricultural Economics Research in A.R.E. (2012).
- A.O.A.C. (1985): Association of Official Agricultural Chemists. Official methods of Analysis, A.O.A.C., Benjamin Franklin Station. Washington D.C., USA.
- Byers, R.E. and C.G. Lyons (1985): Peach flower thinning and possible site of action of desiccating chemicals. J. American Soc. Hort. Sci., 110: 662-667.
- Casanova, L.; R. Cassanova; A. Moret and M. Agusti (2009): The application of gibberellic acid increases berry size of "Emperatriz" Seedless grape. Spanish J. of Agric. Res., 7 (4): 919-927.
- Colapietra, M.; L. Tarricone and G. Tagliente (1995): Effects of gibberellic acid and cluster thinning on the qualitative characteristics of table grape Centennial Seedless. Rivista di frutticoltura ed ortofloricoltura 57 (5): 65-70. (Hort. Abst., 65: 6923).
- Dhillon, W.S.; A.S. Bindra; S.S. Chefma and S. Sahan (1992): Note on effect of berry thinning on quality of grapes cv. Perlette. Indian J. of Hort. 49 (1): 50-52.
- Dokoozlian, N.K. and W.L. Peacock (2001): Gibberellic acid applied at bloom reduces fruit set and improves size of "Crimson Seedless" table grapes. Hort. Sci., Vol. 36, No. 4, pp. 706-709.
- El-Akkad, M.M. (2004): Physiological studies on vegetative growth and fruit quality in some grapevine cultivars. Ph.D. Thesis, Fac. of Agric., Assiut Univ., Egypt, pp. 262.

- El-Hammady, A.M.; A.D. Shaltout; N. Abdel-Hamid and A.M.H. El-Kereamy (1998): Effect of GA<sub>3</sub> treatments on fruit quality of "Flame Seedless" grape cv. Arab Univ. J. Agric. Sci., Ain Shams Univ., Cairo, 6 (2): 531-542.
- El-Moursy, F.M.; F.F. Ahmed and F.A. Abdel-Aziz (1993): Benefits of spraying urea and boron for Red roomy grapevines. Minia First Conf. for Hort. Crop.
- El-Salhy, A.M.; K.I. Ahmed-Amin; A.A.B. Masoud and A.A. Abozeed (2009): Effect of berry thinning, CPPU spraying and pinching on cluster and berry quality of two grapevine cultivars. Assiut J. of Agric. Sci., 40 (4): 92-107.
- El-Salhy, A.M.; K.I. Ahmed-Amin; A.A.B. Masoud and A.A. Abozeed (2011): Response of Ruby Seedless and Red Roomy grapevines to application of some biofertilizers. Assiut J. of Agric. Sci., 41 (5): 125-142.
- Ezzahouani, A.; A.M. Lasheen and L. Walali (1985): Effects of gibberellic acid and girdling on Thompson Seedless and Ruby Seedless table grapes in Morocco. Hort. Sci. 20: 393-394.
- Fawzi, M.I.f.; Laila F. Haggag; M.F. Shahin; M.A. Merwad and E.A. Genaidy (2014): Influence of spraying urea, boron and active dry yeast on growth, yield, leaf chemical composition and fruit quality of Superior grapevines growth in sandy soil conditions. Middle East J. of Applied Sci. 4 (3): 740-747.
- Gomez, K.A. and A.A. Gomez (1984): Statistical Procedures for Agricultural Research 2<sup>nd</sup> Ed. Wiley, New York.
- Guirguis, N.S.; A.H. Gumana; R.G. Stino and A.M. Merhrecki (1996): Effect of carbonate, urea and cyanamide on thinning and apical blooms under arid conditions in Egypt. Hort. Abst. 66: 9200.
- Hassan, A.S.A. (2002): Effect of some GA<sub>3</sub>, yeast, nitrogen, and potassium foliar spray treatments on yield, fruit quality and leaf characteristics of Thompson Seedless. Zagazig J. Agric. Res., Vol. 29 (1): 73-97.
- Idso, S.B.; K.E. Idso, R.L. Garcia; B.A. Kimball and J.K. Hooper (1995): Effect of atmospheric CO<sub>2</sub> enrichment and foliar methanol application on net photosynthesis of sour orange trees (*Citrus aurantium*, Rutaceae) leaves. Amer. J. of Botany, 82 (1): 26-30.
- Lu, J.; O. Lamikanra and S. Leong (1995): Effect of gibberellic acid on muscadine grape production. Proc. Folia. State Hort. Soc. 108: 360-361.
- Markham, K.P. (1982): Techniques of flavonoids identification. Academic Press, London.
- Marzouk, H.A. and H.A. Kassem (2002): Yield and fruit quality at harvest or after storage of Flame Seedless grape as affected by frequent sprays of Gibberellic acid. J. Adv. Agric. Res. Alexandria Univ., 7 (3): 525-537.
- Moor, T.C. (1979): Biochemistry and physiology of plant hormones. Pub. By Springer-Verlag New York, USA.
- Nijjar, G.S. (1985): Nutrition of fruit trees Mrs Usha raj Kumar for Kilyany publishers. New Delhi, India, 206-234
- Omran, Y.A.M. and H.A. Abdel-Latif (2003): Examining of some ge-

- netically improved yeast strains on vine vigor, yield components and fruit quality of Red Roomy grapevines. *Assiut J. Agric. Sci.*, 34 (1): 33-42.
- Omran, Y.A.M.; T. Ramadan and F.M. El-Borai (2005): Application of yeast and/or hydrogen cyanamide for improving vine vigor, bud behavior and fruit quality of Thompson Seedless grapevines. *J. Agric. Sci. Mansoura Univ.*, 30 (7): 4055-4071.
- Orth, C.H.F. (1990): Effect of spraying or dipping Muscat Seedless with gibberellic acid at different flowering stage on berry set and berry size. *Deciduous Fruit Grower* 40 (11): 428-432. (*Hort. Abst.*, 62: 3826).
- Perez, F.J.; C. Vionic and J. Retamales (2000): Bioactive gibberellins in seeded and Seedless grapes; identification and changes in content during berry development. *Am. J. Enol Vitic*, 51: 315-318.
- Poni, S. (2003): Summer pruning in vineyards. *Physiology and cultural aspects. Informatore Agrario*, 59 (26): 37-49.
- Roper, T.R. and L.E. Williams (1989): Net CO<sub>2</sub> assimilation and carbohydrate partitioning of grapevine leaves response to trunk girdling and gibberellic acid application. *Plant Physiology*, 89: 1136-1140.
- Selim, A.A. (2007): Response of Flame Seedless grapes to some improving treatments under Assiut environments. M.Sc. Thesis, Fac. Agric., Assiut Univ., Egypt.
- Snedecor, G.W. and W.G. Cochran (1990): *Statistical Methods* 7<sup>th</sup> Ed. The Iowa State Univ., Press.
- Williams, L.E. and J.E. Ayars (2005): Water use of Thompson Seedless grapevines as affected by the application of gibberellic acid (GA<sub>3</sub>) and trunk girdling-practices to increase berry size. *Agric. and Forest Meteorology*, 129: 85-94.
- Winkler, A.J.; A.J. Cook; W.M. Kliwer and L.A. Lider (1974): *General viticulture*. Published by University of California Press, Barkley.
- Zoffoli, J.P.; B.A. Latorre and P. Naranjo (2009): Preharvest application of growth regulators and their effect on post harvest quality of table grape during cold storage. *Postharvest Biology and Techn-ology*, 51 (2): 183-192.

## تأثير رش حمض الجبريليك واليوريا والخميرة علي إثمار شجيرات العنب الفليم اللابذري تحت ظروف التربة الرملية

النوبي حفني سالم الحلبي<sup>1</sup>، عبد الفتاح مصطفى الصالحي<sup>1</sup>، مؤمن محمد الوصفي<sup>2</sup> و رشاد عبدالوهاب إبراهيم<sup>1</sup>

<sup>1</sup> قسم الفاكهة - كلية الزراعة - جامعة أسيوط

<sup>2</sup> قسم البساتين - كلية الزراعة - جامعة جنوب الوادي - قنا

### المخلص:

أجريت هذه الدراسة خلال ثلاثة مواسم متتالية ٢٠١٠، ٢٠١١، ٢٠١٢ علي شجيرات العنب الفليم اللابذري بمزرعة الكرنك لإنتاج عنب المائدة والتي تقع في محافظة الأقصر - جمهورية مصر العربية. بهدف دراسة تأثير رش حمض الجبريليك واليوريا والخميرة الجافة المنشطة علي المحصول وخصائص العناقيد والحبات. وقد تم رش حمض الجبريليك قبيل التزهير وأثناء اكتمال التزهير وعندما وصل قطر الحبات ٦ مم بينما تم رش اليوريا قبيل التزهير وأثناء اكتمال التزهير وتم رش الخميرة المنشطة عندما وصل قطر الحبات ٦ مم.

ويمكن تلخيص أهم النتائج فيما يلي:

- أدي رش حمض الجبريليك في المراحل الثلاثة إلي زيادة معنوية في طول العنقود والمحصول مع نقص نسبة العقد وعدد الحبات والحبات الصغيرة وبالتالي تحسين صفات العنقود ووزن الحبات.
- سبب رش اليوريا قبيل التزهير وأثناء اكتمال التزهير حدوث زيادة معنوية لطول العنقود مع نقص معنوي في نسبة العقد وعدد الحبات والحبات الصغيرة وبالتالي تحسين صفات العنقود والصفات الكيميائية للحبات.
- أدي رش الخميرة في المرحلة الثالثة (قطر الحبات ٦ مم) عقب رش حمض الجبريليك في مرحلتي (قبل التزهير، اكتمال التزهير) إلي زيادة المحصول وتحسين خصائص الحبات الطبيعية والكيميائية.

من نتائج هذه الدراسة يمكن التوصية برش حمض الجبريليك في المراحل الثلاثة للحصول علي محصول عالي وعناقيد وحبات جيدة - كذلك يمكن استبدال رش حمض الجبريليك برش اليوريا وذلك لاستطالة العنقود وخف الحبات وبالمثل استخدام الخميرة بدلاً من حمض الجبريليك في المرحلة الثالثة وذلك لتلافي أضرار رش الجبريليك خصوصاً علي الأصناف الملونة (الفليم). وبالتالي الحصول علي محصول عال مبكر ذو عناقيد وحبات ذات خصائص ممتازة تتفق مع سوق التصدير والقدرة التنافسية بالأسواق الخارجية.