

(Original Article)



## Quality Properties of African Catfish (*Clarias Gariepinus*) Meat as Affected by its Color

Zeinab M. A. Heikal<sup>1\*</sup>; Safaa A. Limam<sup>2</sup>; Mohamed B. Omer<sup>2</sup>; Badawy M. D. Mostafa<sup>1</sup> and Ahmed H. khalifa<sup>2</sup>

<sup>1</sup> Meat and Fish Technology Research Department, Food Technology Research Institute, Agriculture Research Center, Giza, Egypt.

<sup>2</sup> Food Science and Technology Department, Faculty of Agriculture, Assiut University, Assiut, Egypt.

\*Correspondence: [zeinabmostafa6591@gmail.com](mailto:zeinabmostafa6591@gmail.com)

DOI: 10.21608/AJAS.2024.271766.1338

©Faculty of Agriculture, Assiut University

### Abstract

Nutritional quality (proximate chemical composition, mineral content, amino acid composition, and fatty acid composition), safety quality (some heavy metals content and microbial analysis) as well as sensory quality were investigated for African catfish (*Clarias gariepinus*) meat obtained from black color and gray color catfish. Fillet of catfish constituted 48.15 and 47.41 % of total body weight in black and gray catfish respectively. Catfish meat contained 74.62 and 76.00% protein, 21.17 and 19.81% of fat (on dry wet) in black and gray catfish respectively. Gray color catfish fillet recorded higher contents of K, P, Ca, Na, Mg and Fe compared with the mineral content of black color catfish fillet while, the latter had higher concentrations of Zn, Se and Mn. Essential amino acids (EAAs) to Non-essential amino acids (N-EAAs) ratio ranged from 0.72 to 0.74, while, The polyunsaturated fatty acids/ saturated fatty acids (PUFA/SFA) ratio ranged from 0.89 to 0.93 in catfish meat. The results indicated that, catfish meat had a high nutritional, and safety qualities. Regarding to sensory quality, the panelists relatively preferred the meat of gray color catfish fillet. Moreover, some sensory parameters (color, odor, taste, and overall acceptability) of catfish meat may be need pre-treatments to improve its sensory quality.

**Keywords:** Catfish, Color, Nutritional quality, Safety quality, Sensory quality.

### Introduction

Fish is an important food source in human consumption due to its minerals and fatty acids needed for various body functions (Wijayanti *et al.*, 2023). However, the chemical composition of fish varies greatly from one species and one individual fish to another depending on age, sex, environment and season (FAO, 2001). Catfish (*C. gariepinus*) is most important aquaculture freshwater species with tilapia in Egypt. In 2017, the world aquaculture production quantity of farmed catfish species (*i.e.*, freshwater fishes of the order Siluriformes) is 5.5 million tones as percent of 4.93% (FAO, 2019). Catfish is of the lowest price fish in Egypt compared with other species. Generally, Egyptian consumers prefer the

light color catfish and many of them reject the dark color one. Consumers have a psychological belief that, black color catfish is low nutritional value and more contaminated. So, the present study was conducted to estimate the nutritional value and some quality properties of African black and gray colored catfish.

### Materials and Methods

Twenty kg of live catfish (*Claries gariepinus*) with average weight from 2-3 Kg were purchased from a local market in Assiut Governorate, Egypt during 2022, and transported in icebox directly to the Food Technology Laboratory, Agriculture Research Station (ARC), in Assiut city.

Catfish samples were divided according to its color to gray and black catfish as indicated in figure (1). Fish were slaughtered and left to bleed for three minutes, then cut up to main parts. Fillet was minced by a kitchen meat mincer with a 3 mm diameter perforated plate and packaged in polyethylene bags and kept in frozen at  $-18\pm 1^{\circ}\text{C}$  until use for analysis.



**Fig 1. Black and gray catfish.**

### Nutritional quality

Chemical composition (moisture, crude protein, crude fat and ash contents) was determined according to official methods (AOAC, 2000). Carbohydrates were calculated by difference according to Turhan *et al.*, (2005), while caloric value was calculated as multiply the content of protein and carbohydrates by 4 and by 9 for fat content.

Minerals content was analyzed. Microwave digester (Multiwave GO Plus 50 HZ) was used prior to spectrophotometric analysis of the samples by MPAES4210 (Microwave Plasma -Atomic Emission Spectroscopy) (Agilent, Mulgrave, Victoria, Australia). according to Agilent Technologies, Inc. (2021).

Amino acids were determined chromatography using Beckman Amino Acid Analyzer Model 119CL, according to Pellett and Young (1980). Tryptophan was

determined using spectrophotometric method as described by Sastry and Tummuru (1985). Fatty acids were determined according to Kates (1972) and Rossell *et al.*, (1983).

### **Physico-chemical quality**

To determine pH value, homogenized 10 g of prepared catfish meat fillet, then the sample with 90 ml distilled water for 30 seconds, the pH values were recorded, using a pH-meter (Jenway 3010; Jenway Ltd., Essex, UK) at 20°C, as defined by Fernández-López *et al.*, (2006).

Water holding capacity (WHC) was determined according to Hamm (1960).

$$\text{Water holding capacity (\%)} = \% \text{ Moisture} - \% \text{ Expressible water}$$

### **Color**

Catfish fillet color was measured by chroma meter (Konica Minolta, model CR 410, Japan). Color was expressed using the CIE  $L^*$ ,  $a^*$ , and  $b^*$  color system, lightness  $L^*$  dark (0) to light (100), the redness  $a^*$  values (+) reddish to (-) greenish and the yellowness  $b^*$  values (+) yellowish to (-) bluish were estimated according to Commission International de l'Eclairage, (1978). Samples were analyzed at Cairo University Research Park (CURP)/ Faculty of Agriculture.

Thiobarbituric acid (TBA) values were determined according to the method of Pearson, (1991). The total volatile bases nitrogen (TVB-N) was estimated as described by AOAC (2005). Trimethylamine (TMA) was determined according to Malle and Tao, (1987).

### **Microbiological quality**

The total plate bacterial counts were according to procedures by APHA (1976) and Difco- Manual (1984). Yeast and Mould counts (YMC) were determined using Bacto yeast malt (Y.M) according to the methods described by Difco -Manual (1998).

### **Sensory quality**

Sensory qualities were done by 10 staff members of Agriculture Research Station in Assiut city. A 10- point hedonic scale (1 being dislike very much to 10 being like very much) was used according to Gelman and Benjamin (1989).

### **Statistical analysis**

The data obtained from three replicates were analyzed by ANOVA using the SPSS statistical package program, and differences among the means were compared using the Duncan's Multiple Range test (SPSS, 2020). A significance level of 0.05 was chosen.

## **Results and Discussion**

### **Body composition of black and gray catfish (*Clarias gariepinus*) fillet**

The data presented in Table (1) showed the body composition of the studied catfish. Data indicated that, there were significant ( $p > 0.05$ ) differences in the fillet,

head, frame, viscera, skin and fins between black and gray studied catfish, whereas black catfish had heavier weight for the studied parameters. This difference might be due to the age and feeding conditions as reported by Deng (2018). However, the of catfish fillet constituted 48.15 and 47.41% of total body weight in black and gray catfish color; respectively. Such results are agreed with Abd-Elftah (2021) who reported that, the yield of catfish fillet was 48.41%, and Sobczak *et al.*, (2022) showed that, the fillet yield was shown catfish 49.1%. On other hand Abd El-latif (2021) showed that the fillet obtained counted 44.95% of the total body and Abdel-aal (2001) who reported that, the yield of Nile Karmout fish (*Clarias lezera*) fillet was 47.0%. *Clarias gariepinus* has 42.74 % fillet and 54.83 % inedible sections. From the body weight composition one can concluded that the net percentage of the flesh compared to the total body was good to use such kind of fish to produce a fast fish product such as fish burger or fish fingers (Abd El-latif, 2021).

**Table 1. Body composition of black and gray catfish (*Clarias gariepinus*)**

Component	Blackcatfish		Gray catfish	
	Weight (g)	%	Weight (g)	%
<b>Fillet</b>	1178.00±6.504 <sup>a</sup>	48.15	1157.18±7.549 <sup>b</sup>	47.41
<b>Head</b>	532.78±3.196 <sup>a</sup>	21.78	531.86±3.329 <sup>b</sup>	21.79
<b>Frame</b>	308.97±5.839 <sup>a</sup>	12.63	307.17±1.260 <sup>b</sup>	12.58
<b>Viscera</b>	235.09±5.007 <sup>a</sup>	9.61	234.17±3.746 <sup>b</sup>	9.59
<b>Skin</b>	122.46±1.823 <sup>b</sup>	5.98	141.97±4.677 <sup>a</sup>	5.82
<b>Fins</b>	69.07±1.988 <sup>a</sup>	2.82	68.57±2.735 <sup>b</sup>	2.81
<b>Whole body(n=3)</b>	2446.32±9.065 <sup>a</sup>	100	2440.92±8.401 <sup>b</sup>	100

Different letters in the same row means significantly difference (p<0.05)

### Gross Chemical Composition of black and gray catfish fillet

Proximate chemical composition and caloric value of black and gray catfish are shown in Table (2). Moisture, crude protein and fat content had significant differences (p < 0.05) between the compared studied colors catfish. However, gray catfish fillet showed the greatest moisture and protein content 74.16 and 76.00 %; respectively. The consumption of catfish could help to prevent some problems due to deficiencies of protein in the diet of people in developing countries (FAO, 2008). On other hand, the black catfish fillet showed highest fat content and caloric value (21.17% and 498.70 kcal/100g on dry weight, respectively). Moreover, there were no significant differences for both ash and carbohydrate content between the two colors studied catfish as shown in Table (2). Similar results were reported by Yelouassi *et al.* (2018); Mahboob *et al.* (2019); Abiodun *et al.* (2021); (Mahmoud *et al.* (2021); Hamad (2021), Mostafa *et al.* (2023) and Zaghloola *et al.* (2023) on catfish fillet composition. Moreover, the main components of fish meat are water (66–81%), protein (16–21%), carbohydrates (<0.5%), lipids (0.2–25%), and ash (1.2–1.5%) in wet weight basis as reported by FAO (1999).

The disparity of chemical composition could be attributed to age and feeding conditions as reported by Deng (2018).

**Table 2. Proximate chemical composition of black and gray catfish fillet (dry weight bases)**

Parameters	Black catfish	Gray catfish
Moisture	73.88 ±0.440 <sup>a</sup>	74.16±1.00 <sup>b</sup>
Protein%*	74.62±0.300 <sup>b</sup>	76.00±0.240 <sup>a</sup>
Fat%*	21.17±1.00 <sup>a</sup>	19.81±1.26 <sup>b</sup>
Ash%*	2.99±0.490 <sup>ab</sup>	3.02±0.020 <sup>a</sup>
Carbohydrate %*	1.23±0.100 <sup>a</sup>	1.16±0.040 <sup>ab</sup>
Caloric value (kcal/100g)*	498.70±1.00 <sup>a</sup>	486.93±0.966 <sup>b</sup>

Different letters in the same row means significantly difference ( $p < 0.05$ ),\*On dry weight bases

### Mineral content of black and gray catfish (*Clarias gariepinus*) fillet

The obtained data in Table (3) revealed that there were eleven minerals in both catfish samples, ten elements as nutritional minerals (Ca, Na, K, P, Mg, Fe, Zn, Mg, Cu, and Cr) and three heavy metals (Hg, Cd and Pb). Gray color catfish fillet was higher content of K, P, Ca, Na, Mg and Fe compared with the mineral content of black color catfish fillet. While black color catfish fillet contains higher concentrations of Zn, Se and Mn. Potassium recorded 4970.09 and 5948.92 ppm in black and gray catfish, respectively. However, potassium supports healthy nerve, muscle, and heart, as well as sugar metabolism, acid-base balance, and brain oxygenation (Rasul *et al.*, 2021). Phosphorous was found as 4119.53 and 5265.27 ppm in black and gray catfish, respectively. On the other hand phosphorous aids in a variety of physiological functions, including phospholipid and adenosine polyphosphate activity (Nair and Mathew, 2001). Black catfish meat had high content of zinc (36.36 ppm) than that in gray catfish meat (29.23 ppm). Moreover, zinc supports the immune system's function, growth and development, and fortifies their defenses against attacks by free radicals (Jónsson *et al.*, 2007). Beside, sodium value recorded 783.82 and 880.42 ppm in black and gray catfish meat, respectively. Also, sodium and potassium help in regulating osmotic pressure in the body. However as indicated in Table (3), iron content was 170.07 and 178.98 ppm in black and gray catfish, respectively. Iron helps in transporting oxygen, calcium aids in protecting bone health, and iodine plays a role in controlling normal growth mechanisms, as well as physical and mental development (Prashanth *et al.* 2015; Goff 2018; Alagawany *et al.* 2021). Gray catfish content 479.05 ppm of magnesium while it was 424.87 ppm in black catfish. Mg is an active component of several enzyme systems in which thymine pyrophosphate is a cofactor. Oxidative phosphorylation is greatly reduced in the absence of Mg while Mn is a cofactor of hydrolase, decarboxylase, and transferase enzymes (Murray *et al.*, 2000). Also, gray catfish meat recorded higher content of Cu (2.69 ppm) compared to black catfish meat (1.63 ppm). Copper is a constituent of many enzymes, and plays a role in iron absorption (Chandra, 1990). Chromium content was ranged from 0.20 to 0.22 ppm in catfish meat. Finally, Cr deficiencies may exist, particularly in children suffering from protein-calorie malnutrition (Mertz, 1974). Similar results were reported by Ersoy and Özeren (2009); Adelakun *et al.* (2017); Abd-Elftah (2021); Abiodun *et al.* (2021) and Wijayanti *et al.* (2023) for different prepared catfish fillet.

The variations in mineral values could be as a result of the environmental conditions, feed, ecological requirements, metabolism and other factors such as salinity, water pollution level, and sediment (Ashraf *et al.* 2012). Moreover, the differences in the mineral content could be attributed to the habitat, type of feeding, type of feeding and age (Abd-Elftah, 2021 and Abiodun *et al.*,2021).

**Table 3. Minerals content of black and gray catfish fillet**

Minerals	Black catfish	Gray catfish
	<b>mg /kg (ppm)</b>	
<b>Ca</b>	1017.21	1147.47
<b>Na</b>	783.82	880.42
<b>K</b>	4970.09	5948.92
<b>Mg</b>	424.87	479.05
<b>Cu</b>	1.63	2.69
<b>Fe</b>	170.07	178.98
<b>Zn</b>	36.36	29.23
<b>P</b>	4119.53	5265.27
<b>Se</b>	0.20	0.00
<b>Mn</b>	0.61	2.01
<b>Cr</b>	0.20	0.22
	<b>Micro gram/kg</b>	
<b>Hg</b>	<0.1	<0.1
<b>Cd</b>	<0.1	<0.1
<b>Pb</b>	<0.10	<0.10

On the other hand, the heavy metals (Hg, Cd and Pb) were recorded < 0.1 micro gram/kg in black and gray catfish fillet. The European maximum levels of 0.30 mg/kg for Pb and 0.05mg/kg for Cd (FAO, 2003) maximum levels are 0.02 and 0.05; respectively. So, the determined heavy metals in this study were less than this limits in both of catfish colors. This means that the studied catfish fillet is safe for human consumption and cannot impose health hazard to the consumers.

#### **Amino acid profile of black and gray catfish meat (g/100g protein)**

Protein is an important nutrient for growth and development of humans and its quality depends on the types and composition of its amino acids (Oriolowo *et al.*, 2020). The amino acid profile of black and gray colors catfish fillet is presented in Table (4). The results indicated that, all the essential amino acids, were presented in the studied catfish fillet proved that its high-quality sources of protein for nutrition. In addition, the meat of African catfish can be classified as protein products. However, there was a slight difference between amino acid compositions of the two studied catfish fillets as indicated in Table (4). The total essential amino acids content ranged from 41.04 to 42.19 g/100g protein. This was similar to 42.5% reported for catfish by Osibona *et al.*, (2009). Also, Abd-Elftah, (2021) revealed that, total essential amino acids content was 41.81 g/100g protein. On other hand Zaglol and Eltadawy (2009) found that total essential amino acids in crayfish was 55.703%. The  $\Sigma$ EAA in the investigated catfish fillet was higher than in the FAO/WHO standard of (33.9 g/100 g of protein). The, EAAs cannot be produced

by the human body, thus they must be obtained directly from food (Schaafsma, 2000). Moreover, lysine and leucine content recorded the highest value among the essential amino acid. lysine is required for tissue growth and repair; helpful for maintaining the nitrogen balance in adults and necessary for growth in children. Leucine is also helpful for the remodeling and synthesis of muscle proteins (Junianto *et al.*, 2022).

On other hand, E/NE ratio in black and gray recorded 0.72 to 0.74. This result was lower than 0.82 reported by Ahmad *et al.* (2013) and 0.84 reported by Zaglol, and Eltadawy (2009). In this side Abd-Elftah (2021) revealed that, E/NE ratio in catfish meat was 0.72. Also, Oriolowo *et al.*, (2020) found that the essential to non-essential amino acids ratio was 0.69. Tryptophan recorded 1.08 and 1.09 g/100g protein in black and gray colors catfish fillet: respectively. Tryptophan plays an important biological role in the human body. It takes part in the synthesis of serotonin, which is one of the most important neurotransmitters, which makes.

**Table 4. Amino acid profile of black and gray catfish meat (g/100g protein)**

Amino acid	Black catfish	Gray catfish
Threonine	5.06	5.08
Valine	5.19	5.26
Methionine	2.89	2.93
Isoleucine	4.93	4.94
Leucine	8.73	8.71
Phenylalanine	3.91	3.91
Lysine	10.25	10.27
Tryptophan	1.08	1.09
Total EAAs	41.04	42.19
Histidine	2.12	2.15
Arginine	5.98	5.93
Aspartic Acid	9.85	9.88
Serine	3.80	3.76
Glutamic Acid	15.89	15.91
Proline	3.52	3.56
Alanine	6.79	6.82
Cysteine	0.75	0.75
Tyrosine	3.42	3.19
Glycine	5.03	5.43
Total N EAAs	57.15	57.28
Total amino acid	99.19	99.47
EAA/ N EA	0.72	0.74

EAA/ N EA= Essential amino acids (EAAs) to Non-essential amino acids (N-EAAs) ratio

It extremely necessary to enter the body in depression, insomnia. Such results are in agreement with that reported by Oriolowo *et al.* (2020) and Abd-Elfttah, (2021) for catfish and Zaglol, and Eltadawy, (2009) for crayfish meat. However, the data given in the Table (4) show that the obtained results of black and gray color catfish fillet indicated that, the N-EAAs presented the highest concentration were glutamic ranging from 15.89 to 15.91, respectively. Glutamic acid turns in the body into glutamate that helps nerve cells in the brain send and receive information from other cells. Similar result was reported by Adeyeye (2009); Oriolowo *et al.* (2020) and Abd-Elfttah (2021) for catfish meat.

### **Fatty acid profile of black and gray catfish meat**

The fatty acids profile of total lipids of the studied black and gray colors catfish are presented in Table (5). The data revealed that, the total saturated fatty acids of catfish meat was ranged from 29.11 to 29.94% of fatty acids. The dominant saturated fatty acid in catfish fillet was palmitic acid (C16:0) which recorded 22.88-21.03% while, the total unsaturated fatty acids content was ranged from 69.24 to 71.82% of fatty acids. Oleic fatty acid (C18:1) was the dominant unsaturated fatty acid which recorded 35.78-35.89% of total fatty acids content. Moreover, oleic fatty acid constituted more than a third of total fatty acids of catfish meat. Oleic acid is the most important constitute of plasma free fatty acids and could be reported as an anti-inflammatory fatty acid plying a role in the activation of different pathway of immune competent cells (Carrillo *et al.*, 2012).

As indicated in Table (5), linoleic (18:2) comes in the second order of unsaturated fatty acids content of catfish meat and recorded 19.11 to 19.37% followed by linolenic acid (18:3) with 2.95 to 3.08% of total fatty acids content. However, unsaturated fatty acids, which the body is unable to produce, are essential to human metabolism. They are precursors of a class of hormones known as prostaglandins, which are crucial for both the appropriate healing of inflammatory processes and muscular contractions (Coultae, 1989). Generally fat of catfish meat contains a considerable amount of omega-3 and omega-6 fatty acids, which are crucial for a child's brain development and for lowering blood pressure to prevent stroke (Junianto *et al.*, 2022).

As recommended by WHO (2003) the minimum PUFA/SFA ratio could be 0.4 – 0.5 in foods, this ratio in catfish meat was ranged from 0.89-0.99 which indicated that catfish meat is high quality in this concern. Regarding to catfish color, the gray catfish meat continent a high quantity of unsaturated fatty acid compared to the black one.

However, the slight differences may be due to different factors, such as breed, sex, age, diet, geographical location, environment, may all influence the fatty acid composition (Fernández-López *et al.*, 2006). The obtained results for fatty acid composition are the same line with that reported by Ljubojević *et al.* (2013); Yu *et al.* (2017) for wels catfish (*Silurus glanis*), Chauke *et al.* (2008); Abd-Elfttah (2021); Hamad (2021) and Sobczak *et al.* (2022) for raw catfish *gariepinus* and catfish *heteroclaris*.



**Table 5. Fatty acid profile black and gray catfish meat (as % of total fatty acids)**

Fatty acid	Carbon chain	Black catfish	Gray catfish	Recommended minimum ratio (WHO,2003)
Myristic acid	C14:0	0.79	0.75	
Tetradecenoic	C14:1	0.13	0.11	
Pentadecanoic	C15:0	0.17	0.17	
Palmitic acid	C16:0	22.88	21.03	
Palmitoleic	C16:1	6.09	6.07	
Heptadecanoic acid	C17:0	0.16	0.23	
Cis-10- Heptadecenoic	C17:1	0.24	0.26	
Stearic	C18:0	5.75	5.63	
Oleic acid	C18:1	35.78	35.89	
Linoleic acid	C18:2	19.11	19.37	
$\gamma$ - Linolenic acid	C18:3n6	1.95	1.97	
Linolenic acid	C18:3n3	1.00	1.11	
Stearidonic acid	C18:4	0.26	0.28	
Arachidic acid	C20:0	0.19	0.21	
Cis-11- Eicosenoic acid	C20:1	0.34	0.40	
Eicosadienoic acid	C20:2	0.25	0.27	
Eicosatrienoic acid	C20:3n6	0.51	0.59	
	C20:3n3	0.77	1.1	
Eicosapentaenoic acid (EPA)	C20:5	1.14	1.29	
Behenic	C22:0	0.25	0.24	
Docosatetraenoic acid	C22:4	0.57	0.63	
Cludandonic acid (DPA)	C22:5	0.44	0.49	
Docosahexaenoic acid (DHA)	C22:6n3	0.80	1.71	
Lignoceric acid	C24:0	0.14	0.13	
Non identified fatty acid		0.38	0.42	
<b>Total fatty acid</b>		99.71	99.93	
<b>SFA</b>		29.94	29.11	
<b>USFA</b>		69.24	71.82	
<b>MUSFA</b>		42.58	42.73	
<b>PUSFA</b>		26.63	28.91	
<b>PUFA/SFA ratio</b>		0.89	0.99	0.4 – 0.5

SFA= saturated fatty acids, USFA= unsaturated fatty acids, MUSFA= mono unsaturated fatty acids, PUSFA= Poly unsaturated fatty acids, PUFA/SFA= Poly unsaturated fatty acids: saturated fatty acids,

### Physicochemical properties of black and gray catfish fillet

Some physicochemical properties of black and gray catfish fillet are presented in Table (6). Black catfish meat recorded significantly ( $p \leq 0.05\%$ ) higher values of pH and TBA contents compared with gray color catfish meat, while the later was significantly ( $p \leq 0.05\%$ ) higher of WHC and there were no significant of total volatile bases nitrogen ( $p \leq 0.05\%$ ) of total volatile bases nitrogen TVB-N and trimethylamine (TMA) contents between the two studied catfish meats. However, pH of catfish meat ranged from 6.50 to 6.54. Generally,

the pH value of meat is considered the main quality indicator through processing technologies and storage. (Cawthorn *et al.*, 2018). Since the pH value is not enough index used alone to determine the quality of fish, it can be used as a guideline to control the quality of fish when joined with other quality parameters (Ozogul *et al.*, 2011). Similar results were reported by Mahboob *et al.* (2019); Abd-Elftah (2021) and (Mahmoud *et al.* (2021) for catfish fillet. The obtained results indicated that WHC of gray catfish meat was 71.18% while it was only 70.81% in black catfish meat and that may be due to the relatively high content of protein as indicated in Table (2). However, measuring the ability of the muscle to hold water is one of many ways to describe quality of fish. This is an extensive characteristic for many reasons: From the point of economical view, weight reduction due to water loss is of very importance (Olsson *et al.*, 2003), The accumulation of discharge is unattractive to consumers and loss of water is negatively affected the muscle texture (Abd El-latif, 2021). The obtained results agreement comparatively with that of Paleckaitis *et al.* (2018); Abd-Elftah (2021) and Mahmoud *et al.* (2021) for catfish fillet. According to data in Table (6), the TBA value in both black and gray catfish fillet was 0.85 mg malonaldehyde / kg, while it was 0.80 mg malonaldehyde / kg gray color catfish fillet and that reflect the relatively high fat content in black catfish meat. Moreover, the maximum TBA value, in good quality fish, is 5 mg (MDA/ kg) as reported by Ozogul *et al.* (2011). Besides Gahrue *et al.* (2017) mentioned that fresh catfish contain 3-5 mg MDA/ kg sample) but if the values increased to 5-8 mg (MDA/ kg sample) it limits the product acceptability.

As indicated in Table (6) TVB-N contents was ranged from 7.91 to 7.93 mg/100g sample while TMA was ranged from 2.01 to 2.02 mg/100g. Moreover, the concentration of TVB-N in freshly caught fish is normally between 5 and 20 mg/100g, while 30-35 mg/100g of flesh is generally considered to be the maximum level of acceptability for cold water fish reported by Muhammet and Sevim (2007).

**Table 6. Physicochemical properties of black and gray catfish fillet**

Parameters	Black catfish	Gray catfish
pH value	6.54±0.003 <sup>a</sup>	6.50±0.020 <sup>b</sup>
WHC (%)	70.81±0.810 <sup>b</sup>	71.18±0.080 <sup>a</sup>
TBA mg MDA/ kg	0.85±0.020 <sup>a</sup>	0.80±0.070 <sup>b</sup>
TVB-Nmg/100g	7.93±0.445 <sup>a</sup>	7.91±0.360 <sup>ab</sup>
TMAN mg/100g	2.02±0.100 <sup>a</sup>	2.01±0.050 <sup>a</sup>

Different letters in the same row means significantly difference ( $p < 0.05$ )

Moreover, the acceptability recommended limit (35mg/ 100g of fish flesh) for fish as defined by EOS (2005). While the acceptability recommended rate for TMA values in fish are 10-15 mg/100g of fish as suggested by Lakshmanan (2000). According to the (EOS, 1991), 30 mg/100g sample is the allowed upper limit of TVB-N and 10 mg/100g sample is the allowed upper limit of TMA. The above mentioned results are in agreement with those obtained by Hamad, (2021), who reported that TVBN and TMAN in raw catfish fillets (*Clarias gariepinus*) were 13.40 and 1.02 mg/100g and (Mahmoud *et al.* (2021) showed that, frozen catfish fillet during the storage period up to 12 months, the TVB-N value increase

from 22.02 to 24.27 mg/100g. Moreover, Mohamed (2016) showed that, the total volatile basis nitrogen (TVB.N) and trimethylamine (TMA) value for fresh Catfish were 6.55 and 3.696 mg/100g; respectively. Abd-Elftah, (2021) found that, the TVB-N in catfish meat was  $5.97 \pm 0.58$  mg/100g.

### Color values (L\*, a\*, and b\*) of black and gray colors catfish fillet

The first thing that catches the consumer's attention in seafood is its color, thus color has an inevitable effect on the food products' acceptability and marketability (Li *et al.*, 2013). Pigmentation is highly valued in some fish, such as salmon and trout, and ornamental fish, while in others, such as channel catfish, *Ictalurus punctatus*, it is highly undesirable (Li *et al.*, 2007).

Date in Table (7) indicates the color values (L, a\*, and b\*) of black and gray catfish fillet. Gray catfish fillet was significantly higher than black catfish in L\* value (lightness), which recorded 52.99 and 51.61, respectively. Whereas the obtained result of a\* value (redness) of black catfish revealed that it was more redness compared with gray catfish fillet (21.05 and 19.40; respectively). On other hand, b\* values (yellowness) recorded 17.04 in gray color catfish and 18.92 in black color catfish. From such results, gray catfish meat seemed to be lighter than the meat of black color catfish. These results agree with Sobczak *et al.* (2022) showed that, (L\*) and (WI), as well as lower a\* and b\* parameters mentioned in the fillets of *heteroclarias* compered than *C. gariepinus*.

**Table 7. color values (L\*, a\*, and b\*) of black and gray catfish fillet**

Parameters	L*	a*	b*
<b>Black catfish</b>	51.61±0.815 <sup>b</sup>	21.05 ±0.767 <sup>a</sup>	18.92±0.692 <sup>a</sup>
<b>Gray catfish</b>	52.99±0.605 <sup>a</sup>	19.40±0.423 <sup>b</sup>	17.04±0.416 <sup>b</sup>

Different letters in the same row means significantly difference (p<0.05)

### Microbial evaluation of black and gray catfish fillet

Microbial activity is generally responsible for the deterioration of fish, total bacterial count is used as an acceptability index for fish fillets and its products due to the impact of bacteria on fish spoilage (Mahmoud *et al.*, 2021).

The mean values of the total bacterial count (TPC) and yeast and mould counts (YMC) per g of catfish fillet are presented in Table (8). The total plate count recorded 4.60 log cfu/g in black catfish fillet which it higher than that of the gray catfish fillet (4.47 log cfu/ g). The results come in agreement with International Commission on Microbiological Specification for Foods (ICMSF, 1986) and stated that the maximum recommended bacterial counts for good quality product is  $5 \times 10^5$  (5.7 log<sub>10</sub> cfu/g). Moreover, the maximum limit of 6 (log CFU/g) is acceptable in fresh fish as reported by Zhu *et al.* (2015). As shown in Table (8), yeast and moulds recorded 2.17and 2.08log cfu/g, respectively in black catfish fillet which it higher than that of the gray catfish fillet recorded 2.08 and 1.82log cfu/g. Mostafa *et al.* (2023) showed that, the raw *Pangasius* fish fillets and Indian mackerel had total plate bacterial counts (TPC) of  $3.8 \times 10^4$  and  $2.5 \times 10^3$  CFU/ g; respectively. While yeast and mold counts were  $6.2 \times 10^2$  and  $0.97 \times 10^2$  CFU/g;

respectively. On other hand these results of total bacterial counts are less than the Egyptian standard (ESO, 2005) acceptable limit of fish (105 CFU/g).

**Table 8. Microbial evaluation as log cfu/g of black and gray catfish fillet**

Parameters	TPC	YMC	
		Yeast	Mould
<b>Black catfish</b>	4.60±0.05 <sup>a</sup>	2.17±0.06 <sup>a</sup>	1.95±0.02 <sup>a</sup>
<b>Gray catfish</b>	4.47 ±0.08 <sup>b</sup>	2.08±0.03 <sup>b</sup>	1.82±0.05 <sup>b</sup>

Different letters in the same row means significantly difference ( $p < 0.05$ ), TPC =Total plate count, YMC= yeast and mold counts

### Sensory evaluation of the black and gray fillet

The sensory characteristics of food items are the main keys for consumer's acceptance, fish and fish products are unique food items especially concerning color, odor and texture properties (Mahmoud *et al.*, 2022). According to the means given by the panelists of grilled samples, sensory scores for studied parameters (Table,9) such as taste, flavor, color, texture and overall acceptability revealed that, there were significant differences observed ( $p > 0.05$ ) amongst the catfish fillet for all the studied parameters. However, the panelists relatively preferred the meat of gray catfish fillet. Besides, the scores of panelists for tested parameters were relatively low. So, catfish meat may be needing some pre-treatments to improve its consumers acceptability. This result was in the same line with Sobczak *et al.* (2022), they showed that, the sensory assessment of, *heteroclaris* meat received better notes than *C. gariepinus*.

**Table 9. Sensory evaluation of the black and gray catfish fillet**

Sample Parameter	Black catfish	Gray catfish
<b>Color</b>	6.85±0.150 <sup>b</sup>	7.71±0.31 <sup>a</sup>
<b>Odor</b>	6.46±0.411 <sup>b</sup>	7.80±0.537 <sup>a</sup>
<b>Taste</b>	6.29±0.296 <sup>b</sup>	7.50±0.408 <sup>a</sup>
<b>Texture</b>	7.10±0.409 <sup>b</sup>	7.45±0.471 <sup>a</sup>
<b>Overall acceptability</b>	6.58±0.632 <sup>b</sup>	7.50±0.437 <sup>a</sup>

Different letters in the same row means significantly difference ( $p < 0.05$ )

### Conclusion

From the obtained results, it could be concluded that catfish meat had high nutritional values, good safety quality in the two studied colors catfish, but may be need pre-treatments to improve its sensory quality.

### References

- A.O.A.C (2000). Association of official analytical chemists, 17th Ed. Of A.O.A.C. International. Published by A.O.A.C. international. Maryland, U.S.A., 1250pp.
- A.O.A.C. (2005). Official methods of analysis of the Association of Official Analytical Chemists, International 18<sup>th</sup> edition, In Horwitz, W. (ED), AOAC, Washington (D.C.), 35: 2-36.
- A.P.H.A. (1976). American Public Health Association, Compendium of Method for the Microbiological Examination of food. Speck, M.L. ed., Washington D.C., U.S.A.

- Abd El-Latif, R. H. (2021). Effect of Natural Antioxidant and Packaging Treatment on Stability of Catfish Product Stored Under Frozen Condition. PhD Thesis, Food and Dairy Sci. Dept. Fac. Of Agric. Minia Univ., Egypt.
- Abdel-aal, H. A. (2001). Using antioxidants for extending the shelf life of frozen Nile karmout (*Clarias lezera*) fish mince. *Journal of Aquatic Food Product Technology*. 10 (4): 87-99.
- Abd-Elftah, H. E.A. (2021). Chemical, Technological and Microbiological Studies on Catfish. PhD Thesis, Food and Dairy Sci. Dept. Fac. Of Agric. Minia Univ., Egypt.
- Abiodun, O. A.; Ojo, A.; Kayode, R. M. O.; Edem, V. E.; Shittu, M. O.; Opaleye, Z. A. and Olayinka, T. N. (2021). Chemical and Microbial Properties of Kiln-Smoked Catfish. *Ife Journal of Science*, 23, (2).
- Adelakun, K.M.; Mustapha, M.K.; Amali, R.P. and Mohammed, N. (2017). Seasonal variation in nutritional quality of catfish (*Clarias gariepinus*) from Upper Jebba Basin, Nigeria. *Journal of Nutrition and Food Science*. 7: 5-9.
- Adeyeye, E. I. (2009). Amino acid composition of three species of Nigerian fish: *Clarias anguillaris*, *Oreochromis niloticus* and *Cynoglossus senegalensis*. *Food Chemistry*, 113 (1): 43– 46.
- Agilent Technologies, Inc. (2021). WWW Agilent. com/ chem /4210mp-aes.
- Ahmad, S.M.; Birnin-Yauri, U.A.; Bagudo B.U. and Sahabi D. M. (2013). Comparative analysis on the nutritional values of crayfish and some insects. *African Journal of Food Science and Technology*, 4 (1): 9-12.
- Alagawany, M.; Elnesr, S.S.; Farag, M.R.; Tiwari, R.; Yattoo, M.I.; Karthik, K.; Michalak, I. and Dharma, K. (2021). Nutritional significance of amino acids, vitamins, and minerals as nutraceuticals in poultry production and health-a comprehensive review. *Vet Q* 41 (1): 1-29. DOI: 10.1080/01165176.2020.1857887.
- Ashraf, M.A.; Maah, M.J. and Yusoff, I. (2012). Bioaccumulation of heavy metals in fish species collected from former tin mining catchment. *International Journal of Environmental Research*, 6 (1): 209-218.
- Carrillo, M. C.; Cavia, d. M. and Alonso-Torre, S. (2012). Role of oleic acid in immune system; mechanism of action; a review ;27(4):978-990.
- Cawthorn, D. M.; Fitzhenry, L. B.; Muchenje, V.; Bureš, D.; Kotrba, R. and Hoffman, L. C. (2018). Physical quality attributes of male and female wild fallow deer (*Dama dama*) muscles. *Meat science*, 137: 168-175.
- Chandra, R.K. (1990). Micro-nutrients and immune functions: An overview. *Annal New York Acad. Sci.* 587: 9-16.
- Chauke, E.; Cukrowska, E.; Thaela-Chimuka, M. J.; Chimuka, L.; Nsengimana, H., and Tutu, H. (2008). Fatty acids composition in South African freshwater fish as indicators of food quality. *Water SA*, 34 (1):119-126.
- Commission Internationale de l'Éclairage (CIE). (1978). Official recommendations on uniform colour spaces. Colour difference equations and metric colour terms, Suppl. No. 2. CIE Publication No. 15 Colourimetry. Paris.
- Coultae, T. (1989). *Food. The chemistry of its components*, 6 th edition, the Royal Society of Chemistry, London: Chap. 4.

- Deng O.O. (2018). Evaluation on nutritive value of four commercial fish species in River Nile. *International Journal of Fisheries and Aquatic Studies* 6 (6): 264-267.
- Difco- Manual (1998). Dehydrated culture media and ingredients. 11th edition. Division of Becton Dickinson and Company, Sparks, Maryland, USA.
- Difco-Manual. (1984). Dehydrated Culture Media and Reagents Microbiological and Clinical Laboratory Procedures, Pub-Difco-Lab-Detroit Michigan, USA. 48232, p 1027.
- Egyptian organization for standardization and quality control (2005). E.O.S.Q.C. ESS 1725-1, 2. Sampling plans for fish and fishery products. In: *Micro-organisms in Foods, Sampling for Microbiological Analysis, Principles and Specific Applications Vol 2 International Commission on Micro-biological Specification for Foods (Ed.)*. Toronto, Canada, 92-104.
- EOS, Egyptian Organization for Standardization (1991). Frozen Fish. Egyptian Standard, A. R. E. No. 1796.
- EOS, Egyptian Organization for Standardization (2005). Smoked Fish, Egyptian Organization for Standardization and Quality, A. R. E. No. 288.
- Ersoy, B., and Özeren, A. (2009). The Effect of Cooking Methods on Mineral and Vitamin Contents of African Catfish. *Food Chemistry* 115(2): 419 - 422.
- FAO. (1999). El pescado fresco: Su calidad y cambios de su calidad. Documento técnico de pesca 348. H.H. Huss (ed.). FAO, Roma. 220. p.
- FAO (2001). The composition of fish, produced by Torry Research Station. FAO in partnership with support unit for international fisheries and aquatic research, SIFAR.
- FAO (2003). Report of the FAO Working Group on the Assessment of Small Pelagic Fish off Northwest Africa.
- FAO, (2008). FAO Fisheries and Aquaculture-Chemical Elements of Fish, <http://www.fao.org/fishery/topics/14820/en>.
- FAO. (2019). Top 10 species groups in global aquaculture 2017. Rome, Italy: Food and Agriculture Organization (FAO); 2019.
- Fernández-López, J., Jiménez, S., Sayas-Barberá, E., Sendra, E., and Pérez-Alvarez, J. A. (2006). Quality characteristics of ostrich (*Struthio camelus*) burgers. *Meat science*, 73(2), 295-303.
- Gahruie, H. H.; Hosseini, S. M. H.; Taghavifard, M. H.; Eskandari, M. H.; Golmakani, M. T. and Shad, E., (2017). Lipid oxidation, color changes, and microbiological quality of frozen beef burgers incorporated with shirazi thyme, cinnamon, and rosemary extracts. *Journal of Food Quality*, <https://doi.org/10.1155/2017/6350156>.
- Gelman, A. and Benjamin, E. (1989). Characteristics of mince from Pond-bred silver carp (*Hypophthalmichthys molitrix*) and preliminary experiments on its use in sausage. *J. Sci. Food Agri.*, 47: 225- 241.
- Goff, J.P. (2018). Invited review: Mineral absorption mechanisms, mineral interactions that affect acid-base and antioxidant status, and diet considerations to improve mineral status. *J Dairy Sci* 101 (4): 2763-2813. DOI. 10.3168/jds.2017-13112.

- Hamad, A.M.A. (2021). Changes in chemical composition, fatty acids and sensory quality of fried catfish fillets (*Clariars gariepinus*). GSC Biological and Pharmaceutical Sciences,15(03), 110–115.
- Hamm, R. (1960). Biochemistry of Meat Hydration. C.F: Advanced Food Research (book), P. 10, 355. Academic Press Inc., New York.
- ICMSF (1986). Microorganisms in Foods 2: Sampling for microbiological analysis. Principles and specific applications, 2nd Ed. Oxford: Blackwell Science, pp. 398.
- Jonsson, A.; Finnbogadottir, G.A.; Porkelsson, G.; Magnusson, H.; Reykdal, O. and Arason, S. (2007). Dried fish as health food: Report. Matis Food Research. Innov. Saf., 32 (7): 1–6.
- Junianto, Kristianto, A.D. O. and Yasmin, N. (2022). Catfish Nugget Product Review Article. Global Scientific Journals. (10) 5. <https://www.globalscientificjournal.com>
- Kates, M. (1972): Techniques of Lipidology. Isolation, Analysis and Identification of Lipids. North Holland Publishing Co, Amsterdam.
- Lakshmanan, P.T. (2000). Fish spoilage and quality assessment. In: Iyer, T.S.G., Kandoran, M.K., Thomas, M., Mathew, P.T. (Eds.), Quality Assurance in Seafood Processing. Central Institute of Fisheries Technology and Society of Fisheries Technology, Cochin, India. Journal of Fishery Technology, 45: 25 - 30.
- Li, M.H.; Robinson, E.H.; Oberle, D.F.; Zimba, P.V. (2007). Effects of various dietary carotenoid pigments on fillet appearance and pigment absorption in channel catfish, *Ictalurus punctatus*. J World Aquacul Soc 38: 557- 563.
- Li, Y.; Liu, S.; Cline, D.; Chen S.; Wang, Y. and Bell, L. N. (2013). Chemical Treatments for Reducing the Yellow Discoloration of Channel Catfish (*Ictalurus punctatus*) Fillets. Journal of Food Science. 78. 10.
- Ljubojević, D.M.; Cirković, V. and Dordevićetal. (2013). Fat quality of marketable fresh water is h species in the Republic of Serbia. Czech Journal of Food Sciences ;31(5):445–450.
- Mahboob, S.; Al-Ghanim, K. A.; Al-Balawi, H. F.; Al-Misned, F. and Ahmed, Z. (2019). Study on Assessment of Proximate Composition and Meat Quality of Fresh and Stored *Clarias Gariepinus* and *Cyprinus Carpio*. Brazilian Journal of Biology, 79(4), 651-658.
- Mahmoud, E.A.; Mohamed, E. and Sharaf, A. M. (2021). Quality Assessment of Egyptian Catfish (*Clarias gariepinus*) Fillet During Frozen Storage. J. of Food and Dairy Sci., Mansoura Univ., 12 (10):253 -258.DOI: 10.21608/jfds.2021.97427.1028
- Mahmoud, M.M.; Bahlol, H.E.M.; Abou-Taleb, M.; Ibrahim, M. A.; El-Desouky, A. I. and Sharoba, A. M. A. (2022). Quality Evaluation of Cold Catfish Soaking with Different Herbal Extracts. Annals of Agric. Sci., Moshtohor. 60 (4).
- Malle, P. and Tao, S.H. (1987). Rapid Quantitative Determination of Trimethylamine using Steam Distillation. Journal of Food Protection, 50: 756-760.
- Mertz, W. (1974). Chromium as a dietary essential for man. In: WGHoekstra *et al* eds., Trace Element Metabolism in Animals, 2nd ed. University Park Press, Baltimore.
- Mohamed, R.A.A. (2016). The Application of Volatile Oils in Preservation of Some Fish Products. Ph.D. Thesis, Fac. Agric., Assiut Univ., Assiut, Egypt.

- Mostafa, M.M.; Youssef, K. M.; abouzieid, A. S. and Amin, H. F. (2023). Quality Changes during Frozen Storage of Hot Smoked Fillets and Spreads. *Egyptian Journal of Aquatic Biology & Fisheries*. 27(1): 47 – 67. [www.ejabf.journals.ekb.eg](http://www.ejabf.journals.ekb.eg)
- Muhammet, B and Sevim K., (2007). "Storage properties of three types of fried whiting balls at refrigerated temperatures." *Turkish journal of fisheries and aquatic sciences* 7.1: 65-70.
- Murray, R.K.; Granner, D.K.; Mayes, P.A and Rodwell, V.W. (2000). *Harper's Biochemistry*, 25th Edition, McGraw-Hill, Health Profession Division, USA.
- Nair, P.G.V. and Mathew, S. (2001). *Biochemical composition of fish and shellfish*, Central Institute of Fisheries Technology, Cochin-682029, ICAR.
- Olsson, G. B.; Olsen, R. L. and Ofstad, R. (2003). Post-mortem structural characteristics and water-holding capacity in Atlantic halibut muscle. *LWT-Food Science and Technology*, 36(1): 125-133.
- Oriolowo, O. B.; John, O. J.; Mohammed, U. B. and Joshua, D. (2020). Amino acids profile of catfish, crayfish and larva of edible dung beetle. *Ife Journal of Science*, 22(2), 9-16.
- Osibona, A. O.; Kusemiju, K. and Akande, G. R. (2009). Fatty Acid Composition and Amino Acid Profile of Two Freshwater Species, African Catfish (*Clarias Gariepinus*) and Tilapia (*Tilapia Zillii*). *African Journal of Food, Agriculture, Nutrition and Development*, 9(1), 608-621.
- Ozogul, Y.; Durmuş, M.; Balıkcı, E.; Ozogul, F.; Ayas, D. and Yazgan, H. (2011). The effects of the combination of freezing and the use of natural antioxidant technology on the quality of frozen sardine fillets (*Sardinella aurita*). *International journal of food science & technology*, 46(2), 236-242.
- Paleckaitis M., Buckiūnienė V., Trepėnaitienė R., Klementaviciūtė J., Racevičiūtė-Stupelienė A., šasytė V., Kudlinskienė I. (2018): Effect of different extruded compound feed on african catfish (*Clarias gariepinus*) productivity, meat chemical and technological parameters. *Veterinarija ir Zootechnika*, 76: 66–70.
- Pearson, D. (1991). *The Chemical Analysis of Food National Collage of Technology*, University of Reading. Weybridge, Surrey, T. and A. Churchill.
- Pellett, P. L. and Young, V. R. (1980). *Nutritional Evaluation of Protein Foods*. *Food and Nutrition Bulletin*, (Suppl. 4).
- Prashanth, L.; Kattapagari, K.K.; Chitturi, R.T.; Baddam, V.R.R. and Prasad, L.K. (2015). A review on role of essential trace element and disease. *J Dr NTR Univ Health Sci* 4 (2): 75-85. DOI: 10.4103/2277-8632.158577.
- Rasul, M.G.; Jahan, I.; Yuan, C.; Sharkar, M.S.I.; Bapary, M.A.J.; Baten, M.A. and Shah, A.K.M.A. (2021). Seasonal variation of nutritional constituents of some freshwater and marine fishes of South Asian Countries: A critical review. *Fundamental Applied Agril.*, 6(2): 51-68.
- Rossell, J. B.; King, B.; and Downes, M. J. (1983). Detection of Adulteration. *J. Am. Oil. Chem. Soc.*, 60, 333.
- Sastry, C.S.P., and Tummuru, M.K. (1985). Spectrophotometric Determination of Tryptophan In Proteins. *Jfood, SCI. Technol*, 22, Pp146:147.



- Schaafsma, G. (2000). The protein digestibility–corrected amino acid score. *The Journal of nutrition*, 130(7), 1865S-1867S.
- Sobczak, M.; Panicz, R.; Sadowski, J; Półgęsek, M. and ochowska-Kujawska. J. Z. (2022). Does Production of *Clarias gariepinus* *Heterobranchus longifilis* Hybrids Influence Quality Attributes of Fillets? *Foods*, 11: 2074. <https://doi.org/10.3390/foods11142074>
- SPSS, (2020). SPSS for windows. Release, 20.0., Standard Version, Armonk, NY: IBM Corp.
- Turhan, S.; Sagir, I. and Ustun, N. S. (2005). Utilization of hazelnut pellicle in low-fat beef burgers. *Meat Sci.*, 71:312-316. Pp146:147.
- Wijayanti, F.; Lisdaniyah, A.; Hasanah, M.; Elfidasari, D. (2023). Minerals and fatty acids profile of armored catfish *Pterygoplichthys pardalis* from Ciliwung River, Indonesia. *Nusantara Bioscience* 15: 58-67.
- Yelouassi, C. A. R., Dossou-Yovo, P., Jacquet, N., and Richel, A. (2018). Influence of Salt on the Biochemical Characteristics of Fermented, Salty and Dried Catfish (*Clarias Gariepinus*) in Benin. *Science*, 6(6): 115-122.
- Yu, L. C.; Zzaman, W.; Akanda, M.d. J. H.; Yang, T. A. and Easa A. M. (2017). Influence of Superheated Steam Cooking on Proximate, Fatty Acid Profile, and Amino Acid Composition of Catfish (*Clarias batrachus*) Fillets. *Turk. J. Fish. Aquat. Sci.* 17: 935-943.
- Zaghloola, A.; Tahaa, I. M.; Nagiba A.; Nasra, A.; Elhamamsyb, S. M.; Abdel-Warithc, A. W. A.; Younisc, E. M.; El-Nawasanyd, M. A.; Bauomid, M. A.; El\_Bahlold, A. A.; Daviese, S. J. and Abdelghny, M. F. (2023). Effect of amla and ginger powders on quality criteria of African catfish (*Clarias gariepinus*) fingers. *Brazilian Journal of Biology*. 83: e270808.
- Zaglol, N. F., and Eltadawy, F. (2009): Study on chemical quality and nutrition value of freshwater crayfish (*Procambarus clarkii*). *Journal of the Arabian aquaculture society*, 4(1): 1-18.
- Zhu, Y.; Zhang, K.; Ma, L.; Huo, N.; Yang, H. and Hao, J. (2015). Sensory, Physicochemical, And Microbiological Changes in Vacuum Packed Channel Catfish (*Clarias Lazera*) Patties During Controlled Freezing-Point Storage. *Food Science and Biotechnology*, 24(4): 49-56.

## خصائص جودة اللحم وعلاقتها بلون القرموط الأفريقي

زينب مصطفى احمد هيكل<sup>1</sup>, صفاء عبد الحميد ليمام<sup>2</sup>, محمد بهاء الدين عمر<sup>2</sup>, بدوي محمد درويش مصطفى<sup>1</sup>, احمد حامد خليفه<sup>2</sup>

<sup>1</sup> قسم بحوث تكنولوجيا اللحوم والأسمك، معهد بحوث تكنولوجيا الأغذية، مركز البحوث الزراعية، الجيزة، مصر.  
<sup>2</sup> قسم علوم وتكنولوجيا الأغذية، كلية الزراعة، جامعه اسيوط، اسيوط، مصر.

### الملخص

تم دراسة الجودة التغذوية (محتوى الرطوبة، البروتين، الدهن، الرماد، الكربوهيدرات، العناصر المعدنية، محتوى الأحماض الأمينية، تركيب الأحماض الدهنية) والجودة الصحية (محتوى بعض العناصر الثقيلة والمحتوى الميكروبي) بالإضافة إلى الجودة الحسية للحم القرموط المتحصل عليه من القرموط ذو اللون الرمادي والقروط ذو اللون الأسود. شكل الجزء القابل للأكل (الفيليه) 48.15، 47.41% من إجمالي وزن الجسم في سمك القرموط الأسود والرمادي على التوالي. وقد احتوي لحم سمك القرموط الأسود والرمادي على 74.62 76.00% من بروتين، 21.17، 19.81% من دهن خام (على أساس الوزن الجاف)، في سمك القرموط الأسود والرمادي على التوالي. كما تبين أن لحم القرموط الرمادي يحتوي على نسب أعلى من البوتاسيوم، الفوسفور، الكالسيوم، والصوديوم، والمغنيسيوم والحديد مقارنة بمحتواها المعدني في لحم القرموط ذو اللون الأسود، في حين كان الأخير يحتوي على تركيزات أعلى من الزنك، السيلينيوم والمنغنيز. وتراوحت نسبة الأحماض الأمينية الأساسية (EAAs) إلى الأحماض الأمينية غير الأساسية (N-EAAs) من 0.72 إلى 0.74، بينما تراوحت نسبة الأحماض الدهنية عديدة عدم التشبع إلى الأحماض الدهنية المشبعة (PUFA / SFA) من 0.89 إلى 0.93 في لحم سمك القرموط تحت الدراسة. ومن أوضحت النتائج المتحصل عليها تبين أن لحم كلا النوعين من القراميط ذو خصائص جودة تغذوية وصحية عالية لكليهما. وفيما يتعلق بالجودة الحسية (لون اللحم، الرائحة، الطعم، ودرجة التقبل العام) فإن لحم القرموط يحتاج إلى بعض المعاملات المبدئية لتحسين الجودة الحسية سواء للحم القرموط ذو اللون الرمادي أو الأسود.