

Table(2): Means of weekly abundance of *T. tabaci* individuals/plant, on two onion varieties cultivated in two planting dates and two locations, 2004-2006 seasons.

Location		Assiut			
Month	Date of inspection	No. individuals of <i>T. tabaci</i>			
		Improved Giza 6		Giza 20	
		1 st planting date	2 nd Planting date	1 st planting date	2 nd Planting date
Dec.	10	3.3		2.2	
	17	3.9		2.8	
	24	4.7	0.7	4.9	0.9
	31	7.2	1.0	8.7	1.7
Jan	7	11.2	2.7	12.7	1.7
	14	47.0	10.0	19.5	12.7
	21	27.2	16.5	15.7	16.8
	28	43.2	8.2	18.9	25.0
Feb.	4	64.7	29.0	30.4	24.9
	11	79.0	42.3	26.4	41.4
	18	104.4	62.0	61.2	45.2
	25	186.7	79.4	96.4	107.2
Mar.	4	211.7	217.3	123.7	96.0
	11	281.3	312.0	131.0	164.5
	18	249.7	351.5	177.5	168.4
	25	209.3	336.7	297.7	316.9
Apr.	1	161.0	138.5	98.8	151.9
	8		67.2		185.0
Total		1695.5	1675.0	1128.5	1360.2
Mean		99.7	104.7	66.4	85.0
Location		Arab Alawamer			
Dec.	6	9.7		3.7	
	13	17.3		14.3	
	20	24.7		12.7	
	27	20.3	13.7	21.3	12.3
Jan	3	44.7	12.7	8.7	5.7
	10	79.0	16.7	15.3	10.0
	17	90.7	31.7	51.0	6.3
	24	106.7	12.0	27.0	6.7
	31	179.7	34.3	40.7	21.3
Feb	7	61.7	155.7	27.0	96.3
	14	106.0	73.0	42.3	119.7
	21	110.0	411.3	18.0	165.0
	28	226.3	269.3	41.3	180.0
Mar.	7	147.7	195.7	55.7	237.3
	14	66.7	89.7	144.7	63.7
	21	16.7	27.0	31.0	48.3
	28	7.0	30.0	16.7	41.0
Apr.	4		7.3		15.7
Total		1314.9	1380.4	571.4	1029.3
Mean		82.2	98.6	35.7	73.5